

Astrophysics from the physical point of view

B. V. Vasiliev

The Annotation

The physics of stars requires from its models and theories to be in accordance with totality of existing astrophysical knowledge. It is basis for the star physics. This is the main difference of astrophysics from all other branches of physics, for which foundations are measurement data.

However at an using the progress of a measuring technics, the important measurement data were obtained by astronomers for few last decades. The existence of these data gives a possibility to establish the star physics on the solid base of an agreement of the theory and data of measurements.

To reach the consent of the theory of stars with data of astronomical measurements, it is necessary to abandon to some astrophysical constructions, which are commonly accepted today. At first, necessary to take into account that the interior of the star is formed by electron-nuclear plasma - the electrically polarized medium. So the equation of the balance of plasma in gravitational field must take into account the role of the gravity-induced electric polarization. The taking into account of the gravity-induced electric polarization of plasmas allows to formulate the theory of a star, in which all main parameters - a mass of the star, its temperature, radius and luminosity - are expressed by determined combinations of the world constants, but star individuality is depended of only two parameters - mass and charge numbers of atomic nuclei from which stellar plasma is composed. This theory gives quantitative explanations with satisfactory accuracy for all dependencies, which are measured by astronomers now.

The special attention attracts the stellar mass spectrum. Theoretically the mass of the star can be calculated on a base of balance equations of the intra-stellar plasma. It turns out that, the majority of stars, except most heavy, is composed by plasma with neutron-excess atomic nuclei. The stability of these nuclei is attached by the specifical mechanism of neutronization acting into dense plasma.

Within the framework of this approach, the theories of the periastron rotation of close binary stars and the magnetic field of stars was considered.

It is important to note, that the taking into account of the gravity-induced electric polarization leads to the other conceptual changes. For instance, it rejects the collapse mechanism of stars at the last stage of their evolutions and hereunder puts the question about a mechanism of the "black holes" formation.

Thereby, at taking into account the gravity-induced effects in plasma, we must refuse some models, which are commonly accepted today and seem obvious. The process of a refusing can be painful for a conservative part of astrophysical community. But this refusing is completely justified and absolutely necessary. Only herewith the quantitative explanation can be obtained for all corresponding measurement data without using of a any fitting parameter. The star physics obtains herewith the reliable foundation in the manner of result of these measurements, as a physical science must have.

Abstract

Astrophysics = the star physics was beginning its development without a supporting of measurement data, which could not be obtained then. Still astrophysics exists without this support, although now astronomers collected a lot of valuable information. This is the main difference of astrophysics from all other branches of physics, for which foundations are measurement data. The creation of the theory of stars, which is based on the astronomical measurements data, is one of the main goals of modern astrophysics. Below the principal elements of star physics based on data of astronomical measurements are described. The theoretical description of a hot star interior is obtained. It explains the distribution of stars over their masses, mass-radius-temperature and mass-luminosity dependencies. The theory of the apsidal rotation of binary stars and the spectrum of solar oscillation is considered. All theoretical predictions are in a good agreement with the known measurement data, which confirms the validity of this consideration.

Contents

1	Introduction	6
1.1	That is the basis of astrophysics?	6
1.2	The basic postulate of astrophysics	7
1.3	The Galileo's method	7
1.4	What does the measurement data express?	8
1.5	About a star theory development	9
2	The energy-favorable state of hot dense plasma	11
2.1	The properties of a hot dense plasma	11
2.1.1	A hot plasma and Boltzman's distribution	11
2.1.2	The correction for the Fermi-statistic	12
2.1.3	The correction for correlation of particles	13
2.2	The energy-preferable state	13
2.2.1	The energy-preferable density	13
2.2.2	The energy-preferable temperature	14
2.2.3	Are accepted assumptions correct?	15
3	The gravity induced electric polarization	16
3.1	Plasma cells	16
3.2	The equilibrium of a nucleus	16
3.3	The equilibrium in electron gas subsystem	17
4	The internal structure of a star	18
4.1	The plasma equilibrium in the star core	18
4.2	The main parameters of a star core	18
4.3	The equilibrium inside the star atmosphere	19
4.4	The radial dependence of density and temperature	19
4.5	The mass of the star	20

5	The main parameters of star	21
5.1	The energy of a star	21
5.1.1	The kinetic energy of plasma	21
5.1.2	The potential energy of star plasma	21
5.2	The temperature of a star core	22
5.2.1	The energy of the black radiation	22
5.2.2	The full energy of a star	23
5.3	Main star parameters	23
5.3.1	The star mass	23
5.3.2	Radii of stars	27
5.3.3	The $\mathbb{R}_*/\mathbb{R}_0$ ratio and $\mathbb{R}_0$	28
5.3.4	The temperature of a star surface	28
5.3.5	The comparison with measuring data	28
6	The thermodynamic relations	31
6.1	The thermodynamic relations	31
6.1.1	The c_P/c_V ratio	31
6.1.2	The Poisson's adiabat	32
6.2	The mass-radius ratio	33
6.3	The mass-temperature-luminosity relations	38
6.3.1	The compilation of the results of calculations	41
7	Magnetic fields of stars	43
7.1	Magnetic moments of stars	43
7.2	Magnetic fields of stars	45
8	The apsidal rotation	48
8.1	The apsidal rotation	48
8.2	The equilibrium form of the core	49
8.3	The angular velocity	50
8.4	The comparison with observations	51
9	The solar seismic oscillations	53
9.1	The spectrum of solar oscillations	53
9.2	The sound speed in plasma	57
9.3	The basic elastic oscillation	57
9.4	The low frequency oscillation	59
9.5	The spectrum of solar oscillations	60
10	Addendum I: A mechanism of stabilization nuclei in plasma	61
10.1	Neutron-excess nuclei and the neutronization	61
10.1.1	The electron cloud of plasma cell	61
10.2	The Thomas-Fermi screening	64
10.3	The screening with relativistic electrons	64
10.4	The neutronization	65

11 Addendum II: Other stars	67
11.1 The atomic substance	67
11.1.1 Small bodies	67
11.1.2 Giants	67
11.2 Plasmas	69
11.2.1 Stars	69
11.2.2 Planet	69
11.2.3 The cold relativistic substance	69
11.2.4 Quasars	71
11.2.5 About the cosmic object categorizations	73
11.3 Several words about star evolution	73
11.4 About «black holes»	75
12 The conclusion	76
Bibliography	79

1 Introduction

1.1 That is the basis of astrophysics?

Astrophysics¹ is linked with other physical sciences using of standard physical laws - laws of mechanics, statistical physics and thermodynamics, electrodynamics etc. However at an inevitable similarity of all physical sciences, astrophysics has an unique peculiar "architecture". Its difference has historical roots.

The modern physics begins its formation at last 16 c. - middle 17 c. mainly with works of W.Gilbert and G.Galileo. They introduce into practice the main instrument of the present exact science - the empirical testing of a scientific hypothesis. Until that time false scientific statements weren't afraid of an empirical testing. A flight of fancy was dainty and refined than an ordinary and crude material world. The exact correspondence to a check experiment was not necessary for a philosophical theory, it almost discredited the theory in the experts opinion. The discrepancy of a theory and observations was not confusing at that time. Now the empirical testing of all theoretical hypotheses gets by a generally accepted obligatory method of the exact science. As a result all basic statements of physics are sure established and based on the solid foundation of an agreement with measurement data.

Astrophysicists were applying standard physical laws for description of the interior of distant and mysterious stars, when there was nothing known about them except that they exist. As a result, up to these days, astrophysical models of a star interior are elaborated without some hope that they can be checked by comparison with measurement data.

The astrophysical community is thinking that a foundation of their science is not a measuring data, but it is an internal consensus of a totality of astrophysical knowledge, models of stars and their evolution, which gives a sure in an objective character of their adequacy.²

At this basis, the first ideas of "apostles" - A.Eddington, S.Chandrasekhar, G.Bete, K.Schwazshchild and others - play a special "canonical" role. It makes the star physics very conservative. In principle, any new results and ideas must be rejected if they were created by physical science after a "canon" formulating. Such happened with plasma physics. Its main results were obtained in the second half of 20 c. only and they are not included in the foundation of astrophysics despite the fact that stars are composed by plasma.

It is important that at present, the basis of astrophysics can be revised. Using the progress of a measuring technics, astronomers were obtaining during last decades of 20 c. the important measuring data. Now one can find approximately ten such measured dependencies and star parameters. The existence of these data opens a new stage of star physics. Now it can abandon to the

¹The modern astrophysics has a whole series of different branches. It is important to stress that all of them except the physics of hot stars beyond the scope of this consideration; we shall use the term "astrophysics" here and below in its initial meaning - as the physics of stars or more tightly as the physics of hot stars.

²According to words of one of known European astrophysicist (on memories).

star description in pre-Galilean manner and to get up on the solid base of an accord of the theory and data measurement, as it accepted in all other branches of physics.

The comparison of these measuring data and appropriate theoretical models will be considered below in detail.

1.2 The basic postulate of astrophysics

The basic postulate of astrophysics - the Euler equation - was formulated in a mathematical form by L.Euler in a middle of 18th century for the "terrestrial" effects description. This equation determines the equilibrium condition of liquids or gases in a gravitational field:

$$\gamma \mathbf{g} = -\nabla P. \quad (1)$$

According to it the action of a gravity forth $\gamma \mathbf{g}$ (γ is density of substance, $\mathbf{g}$ is the gravity acceleration) in equilibrium is balanced by a forth which is induced by the pressure gradient in the substance. This postulate is surely established and experimentally checked be "terrestrial" physics. It is the base of an operating of series of technical devices - balloons, bathyscaphes and other.

All modern models of stellar interior are obtained on the base of the Euler equation. These models assume that pressure inside a star monotone increases depthward from the star surface. As a star interior substance can be considered as an ideal gas which pressure is proportional to its temperature and density, all astrophysical models predict more or less monotonous increasing of temperature and density of the star substance in the direction of the center of a star.

1.3 The Galileo's method

The correctness of fundamental scientific postulate has to be confirmed by empirical testing. The basic postulate of astrophysics - the Euler equation (Eq.(1)) - was formulated long before the plasma theory has appeared, even long before the discovering of electrons and nuclei. Now we are sure that the star interior must consist of electron-nuclear plasma. The Euler equation describes behavior of liquids or gases - substances, consisting of neutral particles. One can ask oneself: must the electric interaction between plasmas particles be taken into account at a reformulation of the Euler equation, or does it give a small correction only, which can be neglected in most cases, and how is it accepted by modern stellar theories?

To solve the problem of the correct choice of the postulate, one has the Galileo's method. It consists of 3 steps:

- (1) *to postulate a hypothesis about the nature of the phenomenon, which is free from logical contradictions;*
- (2) *on the base of this postulate, using the standard mathematical procedures, to conclude laws of the phenomenon;*
- (3) *by means of empirical method to ensure, that the nature obeys these laws (or not) in reality, and to confirm (or not) the basic hypothesis.*

The use of this method gives a possibility to reject false postulates and theories, provided there exist a necessary observation data, of course.

The modern theory of the star interior is based on the Euler equation Eq.(1). Another possible approach is to take into account all interactions acting in the considered system. Following this approach one takes into account in the equilibrium equation the electric interactions too, because plasma particles have electric charges. An energetic reason speaks that plasma in gravity field must stay electroneutral and surplus free charges must be absent inside it. The gravity action on a neutral plasma may induce a displacements of its particles, i.e. its electric polarization, because electric dipole moments are lowest moments of the analysis of multipoles. If we take into account the electric polarization plasma $\mathfrak{P}$, the equilibrium equation obtains the form:

$$\gamma \mathbf{g} + \frac{4\pi}{3} \mathfrak{P} \nabla \mathfrak{P} + \nabla P = 0, \quad (2)$$

On the basis of this postulate and using standard procedures, the star interior structure can be considered (it is the main substance of this paper).

Below it will be shown that the electric force in a very hot dense plasma can fully compensate the gravity force and the equilibrium equation then obtains the form:

$$\gamma \mathbf{g} + \frac{4\pi}{3} \mathfrak{P} \nabla \mathfrak{P} = 0 \quad (3)$$

and

$$\nabla P = 0, \quad (4)$$

which points to a radical distinction of this approach in comparison to the standard scheme based on Eq.(1).

1.4 What does the the astronomic measurement data express?

Are there actually astronomic measurement data, which can give possibility to distinguish "correct" and "incorrect" postulates? What must one do, if the direct measurement of the star interior construction is impossible?

Are there astronomical measurement data which can clarify this problem? Yes. They are:

1. There is the only method which gives information about the distribution of the matter inside stars. It is the measurement of the periastron rotation of closed double stars. The results of these measurements give a qualitative confirmation that the matter of a star is more dense near to its center.
2. There is the measured distribution of star masses. Usually it does not have any interpretation.
3. The astronomers detect the dependencies of radii of stars and their surface temperatures from their masses. Clear quantitative explanations of these dependencies are unknown.
4. The existence of these dependencies leads to a mass-luminosity dependence. It is indicative that the astrophysical community does not find any

quantitative explanation to this dependence almost 100 years after as it was discovered.

5. The discovering of the solar surface oscillations poses a new problem that needs a quantitative explanation.

6. There are measured data showing that some stars are possessing by the magnetic fields. This data can be analyzed at taking into account of the electric polarization inside these stars.

It seems that all measuring data are listed which can determine the correct choice of the starting postulate.³

It is important to underline that all above-enumerated dependencies are known but they don't have a quantitative (often even qualitative) explanation in frame of the standard theory based on Eq.(1). Usually one does not consider the existence of these data as a possibility to check the theoretical constructions. Often they are ignored.

It will be shown below that all these dependencies obtain a quantitative explanation in the theory based on the postulate Eq.(3), which takes in to account the electric interaction between particles in the stellar plasma. At that all basic measuring parameters of stars - masses, radii, temperatures - can be described by definite ratios of world constants, and it gives a good agreement with measurement data.

1.5 About a star theory development

The correct choice of the substance equilibrium equation is absolute requirement of an development of the star theory which can be in agreement with measuring data.

To simplify a task of formulation of the such theory, we can accept two additional postulates.

A hot star generates an energy into its central region continuously. At the same time this energy radiates from the star surface. This radiation is not in equilibrium relatively stellar substance. It is convenient to consider that the star is existing in its stationary state. It means that the star radiation has not any changing in the time, and at that the star must radiate from its surface as much energy as many it generates into its core. At this condition, the stellar substance is existing in stationary state and time derivatives from any thermodynamical functions which is characterizing for stellar substance are equal to zero:

$$\frac{dX}{dt} = 0. \quad (5)$$

Particularly, the time derivative of the entropy must be equal to zero in this case. I.e. conditions of an existing of each small volume of stellar substance can

³ From this point of view one can consider also the measurement of solar neutrino flux as one of similar phenomenon. But its result can be interpreted ambiguously because there are a bad studied mechanism of their mutual conversation and it seems prematurely to use this measurement for a stellar models checking.

be considered as adiabatic one in spite of the presence of the non-equilibrium radiation. We shall use this simplification in Chapter VI.

The second simplification can be obtained if to suppose that a stationary star reaches the minimum of its energy after milliards years of development. (Herewith we exclude from our consideration stars with "active lifestyle". The interesting problem of the transformation of a star falls out of examination too).

The minimum condition of the star energy gives possibility to determine main parameters of equilibrium stellar substance - its density and temperature.

It is reasonable to start the development of the star theory from this point. So the problem of existing of the energy-favorable density of the stellar substance and its temperature will be considered in the first place in the next Chapter.

2 The energy-favorable state of hot dense plasma

2.1 The properties of a hot dense plasma

2.1.1 A hot plasma and Boltzman's distribution

Free electrons being fermions obey the Fermi-Dirac statistic at low temperatures. At high temperatures, quantum distinctions in behavior of electron gas disappear and it is possible to consider electron gas as the ideal gas which obeys the Boltzmann's statistics. At high temperatures and high densities, all substances transform into electron-nuclear plasma. There are two tendencies in this case. At temperature much higher than the Fermi temperature $T_F = \frac{\mathcal{E}_F}{k}$ (where $\mathcal{E}_F$ is Fermi energy), the role of quantum effects is small. But their role grows with increasing of the pressure and density of an electron gas. When quantum distinctions are small, it is possible to describe the plasma electron gas as a the ideal one. The criterium of Boltzman's statistics applicability

$$T \gg \frac{\mathcal{E}_F}{k}. \quad (6)$$

hold true for a non-relativistic electron gas with density 10^{25} particles in cm^3 at $T \gg 10^6 K$.

At this temperatures, a plasma has energy

$$\mathcal{E} = \frac{3}{2}kTN \quad (7)$$

and its EOS is the ideal gas EOS:

$$P = \frac{NkT}{V} \quad (8)$$

But even at so high temperatures, an electron-nuclear plasma can be considered as ideal gas in the first approximation only. For more accurate description its properties, the specificity of the plasma particle interaction must be taken into account and two main corrections to ideal gas law must be introduced.

The first correction takes into account the quantum character of electrons, which obey the Pauli principle, and cannot occupy levels of energetic distribution which are already occupied by other electrons. This correction must be positive because it leads to an increased gas incompressibility.

Other correction takes into account the correlation of the screening action of charged particles inside dense plasma. It is the so-called correlational correction. Inside a dense plasma, the charged particles screen the fields of other charged particles. It leads to a decreasing of the pressure of charged particles. Accordingly, the correction for the correlation of charged particles must be negative, because it increases the compressibility of electron gas.

2.1.2 The hot plasma energy with taking into account the correction for the Fermi-statistic

The energy of the electron gas in the Boltzmann's case ($kT \gg \mathcal{E}_F$) can be calculated using the expression of the full energy of a non-relativistic Fermi-particle system [9]:

$$\mathcal{E} = \frac{2^{1/2} V m_e^{3/2}}{\pi^2 \hbar^3} \int_0^\infty \frac{\varepsilon^{3/2} d\varepsilon}{e^{(\varepsilon - \mu_e)/kT} + 1}, \quad (9)$$

expanding it in a series. (m_e is electron mass, ε is the energy of electron and μ_e is its chemical potential).

In the Boltzmann's case, $\mu_e < 0$ and $|\mu_e/kT| \gg 1$ and the integrand at $e^{\mu_e/kT} \ll 1$ can be expanded into a series according to powers $e^{\mu_e/kT - \varepsilon/kT}$. If we introduce the notation $z = \frac{\varepsilon}{kT}$ and conserve the two first terms of the series, we obtain

$$\begin{aligned} I &\equiv (kT)^{5/2} \int_0^\infty \frac{z^{3/2} dz}{e^{z - \mu_e/kT} + 1} \approx \\ &\approx (kT)^{5/2} \int_0^\infty z^{3/2} \left(e^{\frac{\mu_e}{kT} - z} - e^{2(\frac{\mu_e}{kT} - z)} + \dots \right) dz \end{aligned} \quad (10)$$

or

$$\begin{aligned} \frac{I}{(kT)^{5/2}} &\approx e^{\frac{\mu_e}{kT}} \Gamma\left(\frac{3}{2} + 1\right) - \frac{1}{2^{5/2}} e^{\frac{2\mu_e}{kT}} \Gamma\left(\frac{3}{2} + 1\right) \approx \\ &\approx \frac{3\sqrt{\pi}}{4} e^{\mu_e/kT} \left(1 - \frac{1}{2^{5/2}} e^{\mu_e/kT} \right). \end{aligned} \quad (11)$$

Thus, the full energy of the hot electron gas is

$$\mathcal{E} \approx \frac{3V}{2} \frac{(kT)^{5/2}}{\sqrt{2}} \left(\frac{m_e}{\pi \hbar^2} \right)^{3/2} \left(e^{\mu_e/kT} - \frac{1}{2^{5/2}} e^{2\mu_e/kT} \right) \quad (12)$$

Using the definition of a chemical potential of ideal gas (of particles with spin=1/2) [9]

$$\mu_e = kT \log \left[\frac{N_e}{2V} \left(\frac{2\pi \hbar^2}{m_e kT} \right)^{3/2} \right] \quad (13)$$

we obtain the full energy of the hot electron gas

$$\mathcal{E}_e \approx \frac{3}{2} kT N_e \left[1 + \frac{\pi^{3/2}}{4} \left(\frac{a_B e^2}{kT} \right)^{3/2} n_e \right], \quad (14)$$

where $a_B = \frac{\hbar^2}{m_e e^2}$ is the Bohr radius.

2.1.3 The correction for correlation of charged particles in a hot plasma

At high temperatures, the plasma particles are uniformly distributed in space. At this limit, the energy of ion-electron interaction tends to zero. Some correlation in space distribution of particles arises as the positively charged particle groups around itself preferably particles with negative charges and vice versa. It is accepted to estimate the energy of this correlation by the method developed by Debye-Hückel for strong electrolytes [9]. The energy of a charged particle inside plasma is equal to $e\varphi$, where e is the charge of a particle, and φ is the electric potential induced by other particles on the considered particle.

This potential inside plasma is determined by the Debye law [9]:

$$\varphi(r) = \frac{e}{r} e^{-\frac{r}{r_D}} \quad (15)$$

where the Debye radius is

$$r_D = \left(\frac{4\pi e^2}{kT} \sum_a n_a Z_a^2 \right)^{-1/2} \quad (16)$$

For small values of ratio $\frac{r}{r_D}$, the potential can be expanded into a series

$$\varphi(r) = \frac{e}{r} - \frac{e}{r_D} + \dots \quad (17)$$

The following terms are converted into zero at $r \rightarrow 0$. The first term of this series is the potential of the considered particle. The second term

$$\mathcal{E} = -e^3 \sqrt{\frac{\pi}{kTV}} \left(\sum_a N_a Z_a^2 \right)^{3/2} \quad (18)$$

is a potential induced by other particles of plasma on the charge under consideration. And so the correlation energy of plasma consisting of N_e electrons and (N_e/Z) nuclei with charge Z in volume V is [9]

$$\mathcal{E}_{corr} = -e^3 \sqrt{\frac{\pi n_e}{kT}} Z^{3/2} N_e \quad (19)$$

2.2 The energy-preferable state of a hot plasma

2.2.1 The energy-preferable density of a hot plasma

Finally, under consideration of both main corrections taking into account the inter-particle interaction, the full energy of plasma is given by

$$\mathcal{E} \approx \frac{3}{2} kT N_e \left[1 + \frac{\pi^{3/2}}{4} \left(\frac{a_B e^2}{kT} \right)^{3/2} n_e - \frac{2\pi^{1/2}}{3} \left(\frac{Z}{kT} \right)^{3/2} e^3 n_e^{1/2} \right] \quad (20)$$

The plasma into a star has a feature. A star generates the energy into its inner region and radiates it from the surface. At the steady state of a star, its substance must be in the equilibrium state with a minimum of its energy. The radiation is not in equilibrium of course and can be considered as a star environment. The equilibrium state of a body in an environment is related to the minimum of the function ([9]20):

$$\mathcal{E} - T_o S + P_o V, \quad (21)$$

where T_o and P_o are the temperature and the pressure of an environment. At taking in to account that the star radiation is going away into vacuum, the two last items can be neglected and one can obtain the equilibrium equation of a star substance as the minimum of its full energy:

$$\frac{d\mathcal{E}_{plasma}}{dn_e} = 0. \quad (22)$$

Now taking into account Eq.(20), one obtains that an equilibrium condition corresponds to the equilibrium density of the electron gas of a hot plasma

$$n_e^{equilibrium} \equiv n_* = \frac{16}{9\pi} \frac{Z^3}{a_B^3} \approx 1.2 \cdot 10^{24} Z^3 cm^{-3}, \quad (23)$$

It gives the electron density $\approx 3 \cdot 10^{25} cm^{-3}$ for the equilibrium state of the hot plasma of helium.

2.2.2 The estimation of temperature of energy-preferable state of a hot stellar plasma

As the steady-state value of the density of a hot non-relativistic plasma is known, we can obtain an energy-preferable temperature of a hot non-relativistic plasma.

The virial theorem [9, 18] claims that the full energy of particles E , if they form a stable system with the Coulomb law interaction, must be equal to their kinetic energy T with a negative sign. Neglecting small corrections at a high temperature, one can write the full energy of a hot dense plasma as

$$\mathcal{E}_{plasma} = U + \frac{3}{2} k T N_e = -\frac{3}{2} k T N_e. \quad (24)$$

Where $U \approx -\frac{GM^2}{\mathbb{R}_0}$ is the potential energy of the system, G is the gravitational constant, M and $\mathbb{R}_0$ are the mass and the radius of the star.

As the plasma temperature is high enough, the energy of the black radiation cannot be neglected. The full energy of the stellar plasma depending on the particle energy and the black radiation energy

$$\mathcal{E}_{total} = -\frac{3}{2} k T N_e + \frac{\pi^2}{15} \left(\frac{kT}{\hbar c} \right)^3 V k T \quad (25)$$

at equilibrium state must be minimal, i.e.

$$\left(\frac{\partial \mathcal{E}_{total}}{\partial T} \right)_{N,V} = 0. \quad (26)$$

This condition at $\frac{N_e}{V} = n_*$ gives a possibility to estimate the temperature of the hot stellar plasma at the steady state:

$$\mathbb{T}_* \approx Z \frac{\hbar c}{k a_B} \approx 10^7 Z \text{ K}. \quad (27)$$

The last obtained estimation can raise doubts. At "terrestrial" conditions, the energy of any substance reduces to a minimum at $T \rightarrow 0$. It is caused by a positivity of a heat capacity of all of substances. But the steady-state energy of star is negative and its absolute value increases with increasing of temperature (Eq.(24)). It is the main property of a star as a thermodynamical object. This effect is a reflection of an influence of the gravitation on a stellar substance and is characterized by a negative effective heat capacity. The own heat capacity of a stellar substance (without gravitation) stays positive. With the increasing of the temperature, the role of the black radiation increases ($\mathcal{E}_{br} \sim T^4$). When its role dominates, the star obtains a positive heat capacity. The energy minimum corresponds to a point between these two branches.

2.2.3 Are accepted assumptions correct?

At expansion in series of the full energy of a Fermi-gas, it was supposed that the condition of applicability of Boltzmann-statistics (6) is valid. The substitution of obtained values of the equilibrium density n_* (23) and equilibrium temperature $\mathbb{T}_*$ (27) shows that the ratio

$$\frac{\mathcal{E}_F(n_*)}{k \mathbb{T}_*} \approx 3.1 Z \alpha \ll 1. \quad (28)$$

Where $\alpha \approx \frac{1}{137}$ is fine structure constant.

At appropriate substitution, the condition of expansion in series of the electric potential (17) gives

$$\frac{r}{r_D} \approx (n_*^{1/3} r_D)^{-1} \approx \alpha^{1/2} \ll 1. \quad (29)$$

Thus, obtained values of steady-state parameters of plasma are in full agreement with assumptions which was made above.

3 The gravity induced electric polarization in a dense hot plasma

3.1 Plasma cells

The existence of plasma at energy-favorable state with the constant density n_* and the constant temperature $\mathbb{T}_*$ puts a question about equilibrium of this plasma in a gravity field. The Euler equation in commonly accepted form Eq.(1) disclaims a possibility to reach the equilibrium in a gravity field at a constant pressure in the substance: the gravity inevitably must induce a pressure gradient into gravitating matter. To solve this problem, it is necessary to consider the equilibrium of a dense plasma in an gravity field in detail. At zero approximation, at a very high temperature, plasma can be considered as a "jelly", where electrons and nuclei are "smeared" over a volume. At a lower temperature and a high density, when an interpartical interaction cannot be neglected, it is accepted to consider a plasma dividing in cells [12]. Nuclei are placed at centers of these cells, the rest of their volume is filled by electron gas. Its density decreases from the center of a cell to its periphery. Of course, this dividing is not freezed. Under action of heat processes, nuclei move. But having a small mass, electrons have time to trace this moving and to form a permanent electron cloud around nucleus, i.e. to form a cell. So plasma under action of a gravity must be characterized by two equilibrium conditions:

- the condition of an equilibrium of the heavy nucleus inside a plasma cell;
- the condition of an equilibrium of the electron gas, or equilibrium of cells.

3.2 The equilibrium of a nucleus inside plasma cell filled by an electron gas

At the absence of gravity, the negative charge of an electron cloud inside a cell exactly balances the positive charge of the nucleus at its center. Each cell is fully electroneutral. There is no direct interaction between nuclei.

The gravity acts on electrons and nuclei at the same time. Since the mass of nuclei is large, the gravity force applied to them is much larger than the force applied to electrons. On the another hand, as nuclei have no direct interaction, the elastic properties of plasma are depending on the electron gas reaction. Thus er have a situation, when the force applied to nuclei must be balanced by the force of the electron subsystem. The cell obtains an electric dipole moment d_s , and the plasma obtains polarization $\mathfrak{P} = n_s d_s$, where n_s is the density of the cell.

It is known [10], that the polarization of neighboring cells induces in the considered cell the electric field intensity

$$E_s = \frac{4\pi}{3}\mathfrak{P}, \quad (30)$$

and the cell obtains the energy

$$\mathcal{E}_s = \frac{d_s E_s}{2}. \quad (31)$$

The gravity force applied to the nucleus is proportional to its mass Am_p (where A is a mass number of the nucleus, m_p is the proton mass). The cell consists of Z electrons, the gravity force applied to the cell electron gas is proportional to Zm_e (where m_e is the electron mass). The difference of these forces tends to pull apart centers of positive and negative charges and to increase the dipole moment. The electric field E_s resists it. The process obtains equilibrium at the balance of the arising electric force $\nabla \mathcal{E}_s$ and the difference of gravity forces applied to the electron gas and the nucleus:

$$\nabla \left(\frac{2\pi}{3} \frac{\mathfrak{P}^2}{n_s} \right) + (Am_p - Zm_e)\mathbf{g} = 0 \quad (32)$$

Taking into account, that $\mathbf{g} = -\nabla\psi$, we obtain

$$\frac{2\pi}{3} \frac{\mathfrak{P}^2}{n_s} = (Am_p - Zm_e)\psi. \quad (33)$$

Hence,

$$\mathfrak{P}^2 = \frac{3GM_r}{2\pi r} n_e \left(\frac{A}{Z} m_p - m_e \right), \quad (34)$$

where ψ is the potential of the gravitational field, $n_s = \frac{n_e}{Z}$ is the density of cell (nuclei), n_e is the density of the electron gas, M_r is the mass of a star containing inside a sphere with radius r .

3.3 The equilibrium in plasma electron gas subsystem

Non-uniformly polarized matter can be represented by an electric charge distribution with density [10]

$$\tilde{\varrho} = \frac{\text{div} E_s}{4\pi} = \frac{\text{div} \mathfrak{P}}{3}. \quad (35)$$

The full electric charge of cells placed inside the sphere with radius r

$$Q_r = 4\pi \int_0^r \tilde{\varrho} r^2 dr \quad (36)$$

determinants the electric field intensity applied to a cell placed on a distance r from center of a star

$$\tilde{\mathbf{E}} = \frac{Q_r}{r^2} \quad (37)$$

As a result, the action of a non-uniformly polarized environment can be described by the force $\tilde{\varrho}\tilde{\mathbf{E}}$. This force must be taken into account in the formulating of equilibrium equation. It leads to the following form of the Euler equation:

$$\gamma\mathbf{g} + \tilde{\varrho}\tilde{\mathbf{E}} + \nabla P = 0 \quad (38)$$

4 The internal structure of a star

It was shown above that the state with the constant density is energy-favorable for a plasma at a very high temperature. The plasma in the central region of a star can possess by this property. The calculation made below shows that the mass of central region of a star with the constant density - the star core - is equal to 1/2 of the full star mass. Its radius is approximately equal to 1/10 of radius of a star, i.e. the core with high density take approximately 1/1000 part of the full volume of a star. The other half of a stellar matter is distributed over the region placed above the core. It has a relatively small density and it could be called as a star atmosphere.

4.1 The plasma equilibrium in the star core

In this case, the equilibrium condition (Eq.(32)) for steady density plasma is achieved at

$$\mathfrak{P} = \sqrt{G}\gamma_\star r, \quad (39)$$

Here the mass density is $\gamma_\star \approx \frac{A}{Z}m_p n_\star$. The polarized state of the plasma can be described by a state with an electric charge at the density

$$\tilde{\varrho} = \frac{1}{3}div\mathfrak{P} = \sqrt{G}\gamma_\star, \quad (40)$$

and the electric field applied to a cell is

$$\tilde{\mathbf{E}} = \frac{\mathbf{g}}{\sqrt{G}}. \quad (41)$$

As a result, the electric force applied to the cell will fully balance the gravity action

$$\gamma\mathbf{g} + \tilde{\varrho}\tilde{\mathbf{E}} = 0 \quad (42)$$

at the zero pressure gradient

$$\nabla P = 0. \quad (43)$$

4.2 The main parameters of a star core (in order of values)

At known density $n_\star$ of plasma into a core and its equilibrium temperature $T_\star$, it is possible to estimate the mass $M_\star$ of a star core and its radius $R_\star$. In accordance with the virial theorem⁴, the kinetic energy of particles composing the steady system must be approximately equal to its potential energy with opposite sign:

$$\frac{GM_\star^2}{R_\star} \approx kT_\star N_\star. \quad (44)$$

Where $N_\star = \frac{4\pi}{3}R_\star^3 n_\star$ is full number of particle into the star core.

⁴Below we shall use this theorem in its more exact formulation.

With using determinations derived above (23) and (27) derived before, we obtain

$$\mathbb{M}_\star \approx \frac{\mathbb{M}_{Ch}}{(A/Z)^2} \quad (45)$$

where $\mathbb{M}_{Ch} = \left(\frac{\hbar c}{Gm_p^2}\right)^{3/2} m_p$ is the Chandrasekhar mass.

The radius of the core is approximately equal

$$\mathbb{R}_\star \approx \left(\frac{\hbar c}{Gm_p^2}\right)^{1/2} \frac{a_B}{Z(A/Z)}. \quad (46)$$

where A and Z are the mass and the charge number of atomic nuclei the plasma consisting of.

4.3 The equilibrium state of the plasma inside the star atmosphere

The star core is characterized by the constant mass density, the charge density, the temperature and the pressure. At a temperature typical for a star core, the plasma can be considered as ideal gas, as interactions between its particles are small in comparison with $kT_\star$. In atmosphere, near surface of a star, the temperature is approximately by $3 \div 4$ orders smaller. But the plasma density is lower. Accordingly, interparticle interaction is lower too and we can continue to consider this plasma as ideal gas.

In the absence of the gravitation, the equilibrium state of ideal gas in some volume comes with the pressure equalization, i.e. with the equalization of its temperature T and its density n . This equilibrium state is characterized by the equalization of the chemical potential of the gas μ (Eq.(13)).

4.4 The radial dependence of density and temperature of substance inside a star atmosphere

For the equilibrium system, where different parts have different temperatures, the following relation of the chemical potential of particles to its temperature holds ([9],25):

$$\frac{\mu}{kT} = const \quad (47)$$

As thermodynamic (statistical) part of chemical potential of monoatomic ideal gas is [9],45:

$$\mu_T = kT \ln \left[\frac{n}{2} \left(\frac{2\pi\hbar^2}{mkT} \right)^{3/2} \right], \quad (48)$$

we can conclude that at the equilibrium

$$n \sim T^{3/2}. \quad (49)$$

In external fields the chemical potential of a gas [9]25 is equal to

$$\mu = \mu_T + \mathcal{E}^{potential} \quad (50)$$

where $\mathcal{E}^{potential}$ is the potential energy of particles in the external field. Therefore in addition to fulfillment of condition Eq. (49), in a field with Coulomb potential, the equilibrium needs a fulfillment of the condition

$$-\frac{GM_r\gamma}{rkT_r} + \frac{\mathfrak{P}_r^2}{2kT_r} = const \quad (51)$$

(where m is the particle mass, M_r is the mass of a star inside a sphere with radius r , $\mathfrak{P}_r$ and T_r are the polarization and the temperature on its surface. As on the core surface, the left part of Eq.(51) vanishes, in the atmosphere

$$M_r \sim rkT_r. \quad (52)$$

Supposing that a decreasing of temperature inside the atmosphere is a power function with the exponent x , its value on a radius r can be written as

$$T_r = T_\star \left(\frac{\mathbb{R}_\star}{r} \right)^x \quad (53)$$

and in accordance with (49), the density

$$n_r = n_\star \left(\frac{\mathbb{R}_\star}{r} \right)^{3x/2}. \quad (54)$$

Setting the powers of r in the right and the left parts of the condition (52) equal, one can obtain $x = 4$.

Thus, at using power dependencies for the description of radial dependencies of density and temperature, we obtain

$$n_r = n_\star \left(\frac{\mathbb{R}_\star}{r} \right)^6 \quad (55)$$

and

$$T_r = T_\star \left(\frac{\mathbb{R}_\star}{r} \right)^4. \quad (56)$$

4.5 The mass of the star atmosphere and the full mass of a star

After integration of (55), we can obtain the mass of the star atmosphere

$$\mathbb{M}_A = 4\pi \int_{\mathbb{R}_\star}^{\mathbb{R}_0} (A/Z)m_p n_\star \left(\frac{\mathbb{R}_\star}{r} \right)^6 r^2 dr = \frac{4\pi}{3} (A/Z)m_p n_\star \mathbb{R}_\star^3 \left[1 - \left(\frac{\mathbb{R}_\star}{\mathbb{R}_0} \right)^3 \right] \quad (57)$$

It is equal to its core mass (to $\frac{\mathbb{R}_\star^3}{\mathbb{R}_0^3} \approx 10^{-3}$), where $\mathbb{R}_0$ is radius of a star.

Thus, the full mass of a star

$$\mathbb{M} = \mathbb{M}_A + \mathbb{M}_\star \approx 2\mathbb{M}_\star \quad (58)$$

5 The virial theorem and main parameters of a star

5.1 The energy of a star

The virial theorem [9, 18] is applicable to a system of particles if they have a finite moving into a volume V . If their interaction obeys to the Coulomb's law, their potential energy $\mathcal{E}^{potential}$, their kinetic energy $\mathcal{E}^{kinetic}$ and pressure P are in the ratio:

$$2\mathcal{E}^{kinetic} + \mathcal{E}^{potential} = 3PV. \quad (59)$$

On the star surface, the pressure is absent and for the particle system as a whole:

$$2\mathcal{E}^{kinetic} = -\mathcal{E}^{potential} \quad (60)$$

and the full energy of plasma particles into a star

$$\mathcal{E}(plasma) = \mathcal{E}^{kinetic} + \mathcal{E}^{potential} = -\mathcal{E}^{kinetic}. \quad (61)$$

Let us calculate the separate items composing the full energy of a star.

5.1.1 The kinetic energy of plasma

The kinetic energy of plasma into a core:

$$\mathcal{E}_\star^{kinetic} = \frac{3}{2}k\mathbb{T}_\star\mathbb{N}_\star. \quad (62)$$

The kinetic energy of atmosphere:

$$\mathcal{E}_a^{kinetic} = 4\pi \int_{\mathbb{R}_\star}^{\mathbb{R}_0} \frac{3}{2}k\mathbb{T}_\star n_\star \left(\frac{\mathbb{R}_\star}{r}\right)^{10} r^2 dr \approx \frac{3}{7} \left(\frac{3}{2}k\mathbb{T}_\star\mathbb{N}_\star\right) \quad (63)$$

The total kinetic energy of plasma particles

$$\mathcal{E}^{kinetic} = \mathcal{E}_\star^{kinetic} + \mathcal{E}_a^{kinetic} = \frac{15}{7}k\mathbb{T}_\star\mathbb{N}_\star \quad (64)$$

5.1.2 The potential energy of star plasma

Inside a star core, the gravity force is balanced by the force of electric nature. Correspondingly, the energy of electric polarization can be considered as balanced by the gravitational energy of plasma. As a result, the potential energy of a core can be considered as equal to zero.

In a star atmosphere, this balance is absent.

The gravitational energy of an atmosphere

$$\mathcal{E}_a^G = -4\pi G\mathbb{M}_\star \frac{A}{Z} m_p n_\star \int_{\mathbb{R}_\star}^{\mathbb{R}_0} \frac{1}{2} \left[2 - \left(\frac{\mathbb{R}_\star}{r}\right)^3 \right] \left(\frac{\mathbb{R}_\star}{r}\right)^6 r dr \quad (65)$$

or

$$\mathcal{E}_a^G = \frac{3}{2} \left(\frac{1}{7} - \frac{1}{2} \right) \frac{GM_\star^2}{\mathbb{R}_\star} = -\frac{15}{28} \frac{GM_\star^2}{\mathbb{R}_\star} \quad (66)$$

The electric energy of atmosphere is

$$\mathcal{E}_a^E = -4\pi \int_{\mathbb{R}_\star}^{R_0} \frac{1}{2} \varrho \varphi r^2 dr, \quad (67)$$

where

$$\tilde{\varrho} = \frac{1}{3r^2} \frac{d\mathfrak{P}r^2}{dr} \quad (68)$$

and

$$\tilde{\varphi} = \frac{4\pi}{3} \mathfrak{P}r. \quad (69)$$

The electric energy:

$$\mathcal{E}_a^E = -\frac{3}{28} \frac{GM_\star^2}{\mathbb{R}_\star}, \quad (70)$$

and total potential energy of atmosphere:

$$\mathcal{E}_a^{potential} = \mathcal{E}_a^G + \mathcal{E}_a^E = -\frac{9}{14} \frac{GM_\star^2}{\mathbb{R}_\star}. \quad (71)$$

The equilibrium in a star depends both on plasma energy and energy of radiation.

5.2 The temperature of a star core

5.2.1 The energy of the black radiation

The energy of black radiation inside a star core is

$$\mathcal{E}_\star(br) = \frac{\pi^2}{15} kT_\star \left(\frac{kT_\star}{\hbar c} \right)^3 \mathbb{V}_\star. \quad (72)$$

The energy of black radiation inside a star atmosphere is

$$\mathcal{E}_a(br) = 4\pi \int_{R_\star}^{R_0} \frac{\pi^2}{15} kT_\star \left(\frac{kT_\star}{\hbar c} \right)^3 \left(\frac{\mathbb{R}_\star}{r} \right)^{16} r^2 dr = \frac{3}{13} \frac{\pi^2}{15} kT_\star \left(\frac{kT_\star}{\hbar c} \right)^3 \mathbb{V}_\star. \quad (73)$$

The total energy of black radiation inside a star is

$$\mathcal{E}(br) = \mathcal{E}_\star(br) + \mathcal{E}_a(br) = \frac{16}{13} \frac{\pi^2}{15} kT_\star \left(\frac{kT_\star}{\hbar c} \right)^3 \mathbb{V}_\star = 1.23 \frac{\pi^2}{15} kT_\star \left(\frac{kT_\star}{\hbar c} \right)^3 \mathbb{V}_\star \quad (74)$$

5.2.2 The full energy of a star

In accordance with (61), the full energy of a star

$$\mathcal{E}^{star} = -\mathcal{E}^{kinetic} + \mathcal{E}(br) \quad (75)$$

i.e.

$$\mathcal{E}^{star} = -\frac{15}{7}k\mathbb{T}_*\mathbb{N}_* + \frac{16}{13} \frac{\pi^2}{15} k\mathbb{T}_* \left(\frac{k\mathbb{T}_*}{\hbar c} \right)^3 \mathbb{V}_*. \quad (76)$$

The steady state of a star is determined by a minimum of its full energy:

$$\left(\frac{d\mathcal{E}^{star}}{d\mathbb{T}_*} \right)_{\mathbb{N}=const, \mathbb{V}=const} = 0, \quad (77)$$

it corresponds to the condition:

$$-\frac{15}{7}\mathbb{N}_* + \frac{64\pi^2}{13 \cdot 15} \left(\frac{k\mathbb{T}_*}{\hbar c} \right)^3 \mathbb{V}_* = 0. \quad (78)$$

Together with Eq.(23) it defines the equilibrium temperature of a star core:

$$\mathbb{T}_* = \left(\frac{25 \cdot 13}{28\pi^4} \right)^{1/3} \left(\frac{\hbar c}{ka_B} \right) Z \approx Z \cdot 2.13 \cdot 10^7 K \quad (79)$$

5.3 Main star parameters

5.3.1 The star mass

The virial theorem connect kinetic energy of a system with its potential energy. In accordance with Eqs.(71) and (64)

$$\frac{9}{14} \frac{G\mathbb{M}_*^2}{\mathbb{R}_*} = \frac{30}{7} k\mathbb{T}_*\mathbb{N}_*. \quad (80)$$

Introducing the non-dimensional parameter

$$\eta = \frac{G\mathbb{M}_* \frac{A}{Z} m_p}{\mathbb{R}_* k\mathbb{T}_*}, \quad (81)$$

we obtain

$$\eta = \frac{20}{3} = 6.67, \quad (82)$$

and at taking into account Eqs.(23) and (79), the core mass is

$$\mathbb{M}_* = \left[\frac{20}{3} \left(\frac{25 \cdot 13}{28} \right)^{1/3} \frac{3}{4 \cdot 3.14} \right]^{3/2} \frac{\mathbb{M}_{Ch}}{\left(\frac{A}{Z} \right)^2} = 6.84 \frac{\mathbb{M}_{Ch}}{\left(\frac{A}{Z} \right)^2} \quad (83)$$

The obtained equation plays a very important role, because together with Eq.(58), it gives a possibility to predict the total mass of a star:

$$M = 2M_{\star} = \frac{13.68M_{Ch}}{\left(\frac{A}{Z}\right)^2} \approx \frac{25.34M_{\odot}}{\left(\frac{A}{Z}\right)^2}. \quad (84)$$

The comparison of obtained prediction Eq.(84) with measuring data gives a method to check our theory. Although there is no way to determine chemical composition of cores of far stars, some predictions can be made in this way. At first, there must be no stars which masses exceed the mass of the Sun by more than one and a half orders, because it accords to limiting mass of stars consisting from hydrogen with $A/Z = 1$. Secondly, the action of a specific mechanism (see. Sec.10) can make neutron-excess nuclei stable, but it don't give a base to suppose that stars with $A/Z > 10$ (and with mass in hundred times less than hydrogen stars) can exist. Thus, the theory predicts that the whole mass spectrum must be placed in the interval from 0.25 up to approximately 25 solar masses. These predications are verified by measurements quite exactly. The mass distribution of binary stars⁵ is shown in Fig.1 [7].

The mass spectrum of close binary stars⁶ is shown in Fig.2.

The very important element of the direct and clear confirmation of the theory is obviously visible on these figures - both spectra is dropped near the value $A/Z = 1$.

Beside it, one can easy see it, that the mass spectrum of binary stars (Fig.(1)) consists of series of well-isolated lines which are representing the stars with integer values of ratios $A/Z = 3, 4, 5, \dots$, corresponding hydrogen-3,4,5 ... or helium-6,8,10 ... (also line with the half-integer ratio $A/Z = 3/2$, corresponding, probably, to helium-3, Be-6, C-9...). The existence of stable stars with ratios $A/Z \geq 3$ raises questions. It is generally assumed that stars are composed of hydrogen-1, deuterium, helium-4 and other heavier elements with $A/Z \approx 2$. Nuclei with $A/Z \geq 3$ are the neutron-excess and so short-lived, that they can not build a long-lived stars. Neutron-excess nuclei can become stable under the action of mechanism of neutronization, which is acting inside the dwarfs. It is accepted to think that this mechanism must not work into the stars. The consideration of the effecting of the electron gas of a dense plasma on the nucleus is described in chapter (10). The calculations of Ch.(10) show that the electron gas of dense plasma should also lead to the neutronization mechanism and to the stabilization of the neutron-excess nuclei. This explains the existence of a stable of stars, where the plasma consists of nuclei with $A/Z \geq 3$.

Beside it, at considering of Fig.(1), the question is arising: why there are so few stars, which are composed by very stable nuclei of helium-4? At the

⁵The use of these data is caused by the fact that only the measurement of parameters of binary star rotation gives a possibility to determine their masses with satisfactory accuracy.

⁶The data of these measurements were obtained in different observatories of the world. The last time the summary table with these data was gathered by Khaliulilin Kh.F. (Sternberg Astronomical Institute) [8] in his dissertation (in Russian) which has unfortunately has a restricted access. With his consent and for readers convenience, we place that table in Appendix on <http://astro07.narod.ru>.




Figure 1: The mass distribution of binary stars [7]. On abscissa, the logarithm of the star mass over the Sun mass is shown. Solid lines mark masses, which agree with selected values of A/Z from Eq.(84).




Figure 2: The mass distribution of close binary stars [8]. On abscissa, the logarithm of the star mass over the Sun mass is shown. Solid lines mark masses, which agree with selected values of A/Z from Eq.(84). The binary star spectrum is shown for comparison.

same time, there are many stars with $A/Z = 4$, i.e. consisting apparently of a hydrogen-4, as well as stars with $A/Z = 3/2$, which hypothetically could be composed by another isotope of helium - helium-3. This equation is discussed in Ch.(10).

Beside it, it is important to note, that according to Eq.(84) the Sun must consist from a substance with $A/Z = 5$. This conclusion is in a good agreement with results of consideration of solar oscillations (Chapter 9).

5.3.2 Radii of stars

Using Eq.(23) and Eq.(83), we can determine the star core radius:

$$\mathbb{R}_\star = 1.42 \frac{a_B}{Z(A/Z)} \left(\frac{\hbar c}{Gm_p^2} \right)^{1/2} \approx \frac{9.79 \cdot 10^{10}}{Z(A/Z)} \text{cm}. \quad (85)$$

The temperature near the star surface is relatively small. It is approximately by 3 orders smaller than it is inside the core. Because of it at calculation of surface parameters, we must take into consideration effects of this order, i.e. it is necessary to take into account the gravity action on the electron gas. At that it is convenient to consider the plasma cell as some neutral quasi-atom (like the Thomas-Fermi atom). Its electron shell is formed by a cloud of free electrons.

Each such quasi-atom is retained on the star surface by its negative potential energy

$$(\mathcal{E}_{\text{gravitational}} + \mathcal{E}_{\text{electric}}) < 0. \quad (86)$$

The electron cloud of the cell is placed in the volume $\delta V = \frac{4\pi}{3}r_s^3$, (where $r_s \approx \left(\frac{Z}{n_e}\right)^{1/3}$) under pressure P_e . The evaporation of plasma cell releases energy $\mathcal{E}_{PV} = P_e V_s$, and the balance equation takes the form:

$$\mathcal{E}_{\text{gravitational}} + \mathcal{E}_{\text{electric}} + \mathcal{E}_{PV} = 0. \quad (87)$$

In cold plasma, the electron cloud of the cell has energy $\mathcal{E}_{PV} \approx e^2 n_e^{1/3}$. in very hot plasma at $kT \gg \frac{Z^2 e^2}{r_s}$, this energy is equal to $\mathcal{E}_{PV} = \frac{3}{2} Z kT$. On the star surface these energies are approximately equal:

$$\frac{kT_0}{e^2 n_e^{1/3}} \approx \frac{1}{\alpha} \left(\frac{\mathbb{R}_0}{\mathbb{R}_\star} \right)^2 \approx 1. \quad (88)$$

One can show it easily, that in this case

$$\mathcal{E}_{PV} \approx 2Z \sqrt{\frac{3}{2} kT \cdot e^2 n_e^{1/3}}. \quad (89)$$

And if to take into account Eqs.(55)-(56), we obtain

$$\mathcal{E}_{PV} \approx 1.5 Z kT_\star \left(\frac{\mathbb{R}_\star}{\mathbb{R}_0} \right)^3 \sqrt{\alpha \pi} \quad (90)$$

The energy of interaction of a nucleus with its electron cloud does not change at evaporation of the cell and it can be neglected. Thus, for the surface

$$\mathcal{E}_{electric} = \frac{2\pi\mathfrak{P}^2}{3n_s} = \frac{2G\mathbb{M}_\star}{\mathbb{R}_0} (Am_p - Zm_e). \quad (91)$$

The gravitational energy of the cell on the surface

$$\mathcal{E}_{gravitational} = -\frac{2G\mathbb{M}_\star}{\mathbb{R}_0} (Am_p + Zm_e). \quad (92)$$

Thus, the balance condition Eq.(87) on the star surface obtains the form

$$-\frac{4G\mathbb{M}_\star Zm_e}{\mathbb{R}_0} + 1.5Zk\mathbb{T}_\star \left(\frac{\mathbb{R}_\star}{\mathbb{R}_0}\right)^3 \sqrt{\alpha\pi} = 0. \quad (93)$$

5.3.3 The $\mathbb{R}_\star/\mathbb{R}_0$ ratio and $\mathbb{R}_0$

With account of Eq.(56) and Eqs.(82)-(81), we can write

$$\frac{\mathbb{R}_0}{\mathbb{R}_\star} = \left(\frac{\sqrt{\alpha\pi} \frac{A}{Z} m_p}{2\eta m_e} \right)^{1/2} \approx 4.56 \sqrt{\frac{A}{Z}} \quad (94)$$

As the star core radius is known Eq.(85), we can obtain the star surface radius:

$$\mathbb{R}_0 \approx \frac{4.46 \cdot 10^{11}}{Z(A/Z)^{1/2}} cm. \quad (95)$$

5.3.4 The temperature of a star surface

At known Eq.(56) and Eq.(79), we can calculate the temperature of external surface of a star

$$\mathbb{T}_0 = \mathbb{T}_\star \left(\frac{\mathbb{R}_\star}{\mathbb{R}_0} \right)^4 \approx 4.92 \cdot 10^5 \frac{Z}{(A/Z)^2} \quad (96)$$

5.3.5 The comparison with measuring data

The mass spectrum (Fig.1) shows that the Sun consists basically from plasma with $A/Z = 5$. The radius of the Sun and its surface temperature are functions of Z too. This values calculated at $A/Z=5$ and differen Z are shown in Table (5.3.3)

Table (5.3.3)

Z	$\mathbb{R}_\odot, cm$ (calculated (95))	$\mathbb{T}_\odot, K$ (calculated (96))
1	$2.0 \cdot 10^{11}$	1961
2	$1.0 \cdot 10^{11}$	3923
3	$6.65 \cdot 10^{10}$	5885
4	$5.0 \cdot 10^{10}$	7845

One can see that these calculated data have a satisfactory agreement the measured radius of the Sun

$$\mathbb{R}_{\odot} = 6.96 \cdot 10^{10} cm \quad (97)$$

and the measured surface temperature

$$\mathbb{T}_{\odot} = 5850 K \quad (98)$$

at $Z = 3$.

The calculation shows that the mass of core of the Sun

$$\mathbb{M}_{\star}(Z = 3, A/Z = 5) \approx 9.68 \cdot 10^{32} g \quad (99)$$

i.t. almost exactly equals to one half of full mass of the Sun

$$\frac{\mathbb{M}_{\star}(Z = 3, A/Z = 5)}{\mathbb{M}_{\odot}} \approx 0.486 \quad (100)$$

in full agreement with Eq.(58).

In addition to obtained determinations of the mass of a star Eq.(84), its temperature Eq.(96) and its radius Eq.(95) give possibility to check the calculation, if we compare these results with measuring data. Really, dependencies measured by astronomers can be described by functions:

$$\mathbb{M} = \frac{Const_1}{(A/Z)^2}, \quad (101)$$

$$\mathbb{R}_0 = \frac{Const_2}{Z(A/Z)^{1/2}}, \quad (102)$$

$$\mathbb{T}_0 = \frac{Const_3 Z}{(A/Z)^2}. \quad (103)$$

If to combine they in the way, to exclude unknown parameter Z , one can obtain relation:

$$\mathbb{T}_0 \mathbb{R}_0 = Const \mathbb{M}^{5/4}, \quad (104)$$

Its accuracy can be checked. For this checking, let us use the measuring data of parameters of masses, temperatures and radii of close binary stars [8]. The results of these measurements are shown in Fig.(3), where the dependence according to Eq.(104). It is not difficult to see that these data are well described by the calculated dependence. It speaks about successfulness of our consideration.

If main parameters of the star are expressed through corresponding solar values $\tau \equiv \frac{\mathbb{T}_0}{\mathbb{T}_{\odot}}, \rho \equiv \frac{\mathbb{R}_0}{\mathbb{R}_{\odot}}$ and $\mu \equiv \frac{\mathbb{M}}{\mathbb{M}_{\odot}}$, that Eq.(104) can be rewritten as

$$\frac{\tau \rho}{\mu^{5/4}} = 1. \quad (105)$$

Numerical values of relations $\frac{\tau \rho}{\mu^{5/4}}$ for close binary stars [8] are shown in the last column of the Table(6.2)(at the end Chapter (6)).




Figure 3: The relation between main parameters of stars (Eq.(104)) and corresponding data of astronomical measurements for close binary stars [8] are shown.

6 The thermodynamic relations of intra-stellar plasma

6.1 The thermodynamic relation of star atmosphere parameters

Hot stars steadily generate energy and radiate it from their surfaces. This is non-equilibrium radiation in relation to a star. But it may be a stationary radiation for a star in steady state. Under this condition, the star substance can be considered as an equilibrium. This condition can be considered as quasi-adiabatic, because the interchange of energy between both subsystems - radiation and substance - is stationary and it does not result in a change of entropy of substance. Therefore at consideration of state of a star atmosphere, one can base it on equilibrium conditions of hot plasma and the ideal gas law for adiabatic condition can be used for it in the first approximation.

It is known, that thermodynamics can help to establish correlation between steady-state parameters of a system. Usually, the thermodynamics considers systems at an equilibrium state with constant temperature, constant particle density and constant pressure over all system. The characteristic feature of the considered system is the existence of equilibrium at the absence of a constant temperature and particle density over atmosphere of a star. To solve this problem, one can introduce averaged pressure

$$\hat{P} \approx \frac{GM^2}{\mathbb{R}_0^4}, \quad (106)$$

averaged temperature

$$\hat{T} = \frac{\int_V T dV}{V} \sim T_0 \left(\frac{\mathbb{R}_0}{\mathbb{R}_*} \right) \quad (107)$$

and averaged particle density

$$\hat{n} \approx \frac{N_A}{\mathbb{R}_0^3} \quad (108)$$

After it by means of thermodynamical methods, one can find relation between parameters of a star.

6.1.1 The c_P/c_V ratio

At a movement of particles according to the theorem of the equidistribution, the energy $kT/2$ falls at each degree of freedom. It gives the heat capacity $c_v = 3/2k$.

According to the virial theorem [9, 18], the full energy of a star should be equal to its kinetic energy (with opposite sign)(Eq.(61)), so as full energy related to one particle

$$\mathcal{E} = -\frac{3}{2}kT \quad (109)$$

In this case the heat capacity at constant volume (per particle over Boltzman's constant k) by definition is

$$c_V = \left(\frac{dE}{dT} \right)_V = -\frac{3}{2} \quad (110)$$

The negative heat capacity of stellar substance is not surprising. It is a known fact and it is discussed in Landau-Lifshitz course [9]. The own heat capacity of each particle of star substance is positive. One obtains the negative heat capacity at taking into account the gravitational interaction between particles.

By definition the heat capacity of an ideal gas particle at permanent pressure [9] is

$$c_P = \left(\frac{dW}{dT} \right)_P, \quad (111)$$

where W is enthalpy of a gas.

As for the ideal gas [9]

$$W - \mathcal{E} = NkT, \quad (112)$$

and the difference between c_P and c_V

$$c_P - c_V = 1. \quad (113)$$

Thus in the case considered, we have

$$c_P = -\frac{1}{2}. \quad (114)$$

Supposing that conditions are close to adiabatic ones, we can use the equation of the Poisson's adiabat.

6.1.2 The Poisson's adiabat

The thermodynamical potential of a system consisting of N molecules of ideal gas at temperature T and pressure P can be written as [9]:

$$\Phi = \text{const} \cdot N + NT \ln P - N c_P T \ln T. \quad (115)$$

The entropy of this system

$$S = \text{const} \cdot N - N \ln P + N c_P \ln T. \quad (116)$$

As at adiabatic process, the entropy remains constant

$$-NT \ln P + N c_P T \ln T = \text{const}, \quad (117)$$

we can write the equation for relation of averaged pressure in a system with its volume (The Poisson's adiabat) [9]:

$$\hat{P} V^{\tilde{\gamma}} = \text{const}, \quad (118)$$

where $\tilde{\gamma} = \frac{c_P}{c_V}$ is the exponent of adiabatic constant. In considered case taking into account of Eqs.(111) and (110), we obtain

$$\tilde{\gamma} = \frac{c_P}{c_V} = \frac{1}{3}. \quad (119)$$

As $V^{1/3} \sim \mathbb{R}_0$, we have for equilibrium condition

$$\hat{P}\mathbb{R}_0 = \text{const.} \quad (120)$$

6.2 The mass-radius ratio

Using Eq.(106) from Eq.(120), we use the equation for dependence of masses of stars on their radii:

$$\frac{\mathbb{M}^2}{\mathbb{R}_0^3} = \text{const} \quad (121)$$

This equation shows the existence of internal constraint of chemical parameters of equilibrium state of a star. Indeed, the substitution of obtained determinations Eq.(95) and (96)) into Eq.(121) gives:

$$Z \sim (A/Z)^{5/6} \quad (122)$$

Simultaneously the observational data of masses, of radii and their temperatures was obtained by astronomers for close binary stars [8]. The dependence of radii of these stars over these masses is shown in Fig.4 on double logarithmic scale. The solid line shows the result of fitting of measurement data $\mathbb{R}_0 \sim \mathbb{M}^{0.68}$. It is close to theoretical dependence $\mathbb{R}_0 \sim \mathbb{M}^{2/3}$ (Eq.121) which is shown by dotted line.

If parameters of the star are expressed through corresponding solar values $\rho \equiv \frac{\mathbb{R}_0}{\mathbb{R}_\odot}$ and $\mu \equiv \frac{\mathbb{M}}{\mathbb{M}_\odot}$, that Eq.(121) can be rewritten as

$$\frac{\rho}{\mu^{2/3}} = 1. \quad (123)$$

Numerical values of relations $\frac{\rho}{\mu^{2/3}}$ for close binary stars [8] are shown in the Table(6.2).




Figure 4: The dependence of radii of stars over the star mass [8]. Here the radius of stars is normalized to the sunny radius, the stars masses are normalized to the mass of the Sum. The data are shown on double logarithmic scale. The solid line shows the result of fitting of measurement data $R_0 \sim M^{0.68}$. The theoretical dependence $R_0 \sim M^{2/3}$ (121) is shown by the dotted line.

The Table(6.2).

The relations of main stellar parameters

N	Star		$\mu \equiv \frac{M}{M_{\odot}}$	$\rho \equiv \frac{R_0}{R_{\odot}}$	$\tau \equiv \frac{T_0}{T_{\odot}}$	$\frac{\rho}{\mu^{2/3}}$	$\frac{\tau}{\mu^{7/12}}$	$\frac{\rho\tau}{\mu^{5/4}}$
1	BW Aqr	1	1.48	1.803	1.043	1.38	0.83	1.15
		2	1.38	2.075	1.026	1.67	0.85	1.42
2	V 889 Aql	1	2.4	2.028	1.692	1.13	1.01	1.15
		2	2.2	1.826	1.607	1.08	1.01	1.09
3	V 539 Ara	1	6.24	4.512	3.043	1.33	1.04	1.39
		2	5.31	4.512	3.043	1.12	1.09	1.23
4	AS Cam	1	3.31	2.58	1.966	1.16	0.98	1.13
		2	2.51	1.912	1.709	1.03	1.0	1.03
5	EM Car	1	22.8	9.35	5.658	1.16	0.91	1.06
		2	21.4	8.348	5.538	1.08	0.93	1.00
6	GL Car	1	13.5	4.998	5.538	0.88	1.08	0.95
		2	13	4.726	4.923	0.85	1.1	0.94
7	QX Car	1	9.27	4.292	4	0.97	1.09	1.06
		2	8.48	4.054	3.829	0.975	1.1	1.07
8	AR Cas	1	6.7	4.591	3.111	1.29	1.02	1.32
		2	1.9	1.808	1.487	1.18	1.02	1.21
9	IT Cas	1	1.4	1.616	1.102	1.29	0.91	1.17
		2	1.4	1.644	1.094	1.31	0.90	1.18
10	OX Cas	1	7.2	4.69	4.068	1.25	1.29	1.62
		2	6.3	4.54	3.93	1.33	1.34	1.79
11	PV Cas	1	2.79	2.264	1.914	1.14	1.05	1.20
		2	2.79	2.264	2.769	1.14	1.05	1.20
12	KT Cen	1	5.3	4.028	2.769	1.32	1.05	1.39
		2	5	3.745	2.701	1.28	1.06	1.35
13	V 346 Cen	1	11.8	8.26	4.05	1.59	0.96	1.53
		2	8.4	4.19	3.83	1.01	1.11	1.12
14	CW Cep	1	11.8	8.263	4.051	1.04	1.06	1.11
		2	11.1	4.954	4.393	1.0	1.08	1.07
15	EK Cep	1	2.02	1.574	1.709	0.98	1.13	1.12
		2	1.12	1.332	1.094	1.23	1.02	1.26
16	α Cr B	1	2.58	3.314	1.555	1.76	0.89	1.57
		2	0.92	0.955	0.923	1.01	0.97	0.98
17	Y Cyg	1	17.5	6.022	5.66	0.89	1.06	0.95
		2	17.3	5.68	5.54	0.85	1.05	0.89

The Table(6.2)(continuation).
The relations of main stellar parameters

N	Star	n	$\mu \equiv \frac{M}{M_{\odot}}$	$\rho \equiv \frac{R_0}{R_{\odot}}$	$\tau \equiv \frac{T_0}{T_{\odot}}$	$\frac{\rho}{\mu^{2/3}}$	$\frac{\tau}{\mu^{7/12}}$	$\frac{\rho\tau}{\mu^{5/4}}$
18	Y 380 Cyg	1	14.3	17.08	3.54	2.89	0.75	2.17
		2	8	4.3	3.69	1.07	1.1	1.18
19	V 453 Cyg	1	14.5	8.607	4.55	1.45	0.95	1.38
		2	11.3	5.41	4.44	1.07	1.08	1.16
20	V 477 Cyg	1	1.79	1.567	1.46	1.06	1.04	1.11
		2	1.35	1.27	1.11	1.04	0.93	0.97
21	V 478 Cyg	1	16.3	7.42	5.09	1.15	1.0	1.15
		2	16.6	7.42	5.09	1.14	0.99	1.13
22	V 541 Cyg	1	2.69	2.013	1.86	1.04	1.05	1.09
		2	2.6	1.9	1.85	1.0	1.6	1.06
23	V 1143 Cyg	1	1.39	1.44	1.11	1.16	0.92	0.92
		2	1.35	1.23	1.09	1.0	0.91	0.92
24	V 1765 Cyg	1	23.5	19.96	4.39	2.43	0.67	1.69
		2	11.7	6.52	4.29	1.26	1.02	1.29
25	DI Her	1	5.15	2.48	2.91	0.83	1.12	0.93
		2	4.52	2.69	2.58	0.98	1.07	1.05
26	HS Her	1	4.25	2.71	2.61	1.03	1.12	1.16
		2	1.49	1.48	1.32	1.14	1.04	1.19
27	CO Lac	1	3.13	2.53	1.95	1.18	1.00	1.12
		2	2.75	2.13	1.86	1.08	1.01	1.09
28	GG Lup	1	6.24	4.12	2.64	1.03	1.08	1.11
		2	2.51	1.92	1.79	1.04	1.05	1.09
29	RU Mon	1	3.6	2.55	2.20	1.09	1.04	1.14
		2	3.33	2.29	2.15	1.03	1.07	1.10
30	GN Nor	1	2.5	4.59	1.33	2.49	0.78	1.95
		2	2.5	4.59	1.33	2.49	0.78	1.95
31	U Oph	1	5.02	3.31	2.80	1.13	1.09	1.23
		2	4.52	3.11	2.60	1.14	1.08	1.23
32	V 451 Oph	1	2.77	2.54	1.86	1.29	1.03	1.32
		2	2.35	1.86	1.67	1.05	1.02	1.07
33	β Ori	1	19.8	14.16	4.55	1.93	0.80	1.54
		2	7.5	8.07	3.04	2.11	0.94	1.98
34	FT Ori	1	2.5	1.89	1.81	1.03	1.06	1.09
		2	2.3	1.80	1.62	1.03	1.0	1.03

The Table(6.2)(continuation).
The relations of main stellar parameters

N	Star	n	$\mu \equiv \frac{M}{M_{\odot}}$	$\rho \equiv \frac{R_0}{R_{\odot}}$	$\tau \equiv \frac{T_0}{T_{\odot}}$	$\frac{\rho}{\mu^{2/3}}$	$\frac{\tau}{\mu^{7/12}}$	$\frac{\rho\tau}{\mu^{5/4}}$
35	AG Per	1	5.36	3.0	2.91	0.98	1.09	1.06
		2	4.9	2.61	2.91	0.90	1.15	1.04
36	IQ Per	1	3.51	2.44	2.27	1.06	1.09	1.16
		2	1.73	1.50	2.27	1.04	1.00	1.05
37	ς Phe	1	3.93	2.85	2.41	1.14	1.08	1.24
		2	2.55	1.85	1.79	0.99	1.04	1.03
38	KX Pup	1	2.5	2.33	1.74	1.27	1.02	1.29
		2	1.8	1.59	1.38	1.08	0.98	1.06
39	NO Pup	1	2.88	2.03	1.95	1.00	1.05	1.05
		2	1.5	1.42	1.20	1.08	0.94	1.02
40	VV Pyx	1	2.1	2.17	1.49	1.32	0.96	1.27
		2	2.1	2.17	1.49	1.32	0.96	1.27
41	YY Sgr	1	2.36	2.20	1.59	1.24	0.96	1.19
		2	2.29	1.99	1.59	1.15	0.98	1.12
42	V 523 Sgr	1	2.1	2.67	1.42	1.63	0.92	1.50
		2	1.9	1.84	1.42	1.20	0.98	1.17
43	V 526 Sgr	1	2.11	1.9	1.30	1.15	0.84	0.97
		2	1.66	1.60	1.30	1.14	0.97	1.10
44	V 1647 Sgr	1	2.19	1.83	1.52	1.09	0.96	1.05
		2	1.97	1.67	4.44	1.06	1.02	1.09
45	V 2283 Sgr	1	3.0	1.96	1.67	0.94	0.88	0.83
		2	2.22	1.66	1.67	0.97	1.05	1.02
46	V 760 Sco	1	4.98	3.02	2.70	1.03	1.06	1.09
		2	4.62	2.64	2.70	0.95	1.11	1.05
47	AO Vel	1	3.2	2.62	1.83	1.21	0.93	1.12
		2	2.9	2.95	1.83	1.45	0.98	1.43
48	EO Vel	1	3.21	3.14	1.73	1.44	0.87	1.26
		2	2.77	3.28	1.73	1.66	0.95	1.58
49	α Vir	1	10.8	6.10	3.25	1.66	0.81	1.34
		2	6.8	4.39	3.25	1.22	1.06	1.30
50	DR Vul	1	13.2	4.81	4.79	0.83	1.06	0.91
		2	12.1	4.37	4.79	0.83	1.12	0.93

6.3 The mass-temperature and mass-luminosity relations

Taking into account Eqs.(56), (27) and (46) one can obtain the relation between surface temperature and the radius of a star

$$T_0 \sim R_0^{7/8}, \quad (124)$$

or accounting for (121)

$$T_0 \sim M^{7/12} \quad (125)$$

The dependence of the temperature on the star surface over the star mass of close binary stars [8] is shown in Fig.(5). Here the temperatures of stars are normalized to the sunny surface temperature (5875 C), the stars masses are normalized to the mass of the Sun. The data are shown on double logarithmic scale. The solid line shows the result of fitting of measurement data ($T_0 \sim M^{0.59}$). The theoretical dependence $T_0 \sim M^{7/12}$ (Eq.125) is shown by dotted line.

If parameters of the star are expressed through corresponding solar values $\tau \equiv \frac{T_0}{T_\odot}$ and $\mu \equiv \frac{M}{M_\odot}$, that Eq.(125) can be rewritten as

$$\frac{\tau}{\mu^{7/12}} = 1. \quad (126)$$

Numerical values of relations $\frac{\tau}{\mu^{7/12}}$ for close binary stars [8] are shown in the Table(6.2).

The analysis of these data leads to few conclusions. The averaging over all tabulated stars gives

$$\langle \frac{\tau}{\mu^{7/12}} \rangle = 1.007 \pm 0.07. \quad (127)$$

and we can conclude that the variability of measured data of surface temperatures and stellar masses has statistical character. Secondly, Eq.(126) is valid for all hot stars (exactly for all stars which are gathered in Tab.(6.2)).

The problem with the averaging of $\frac{\rho}{\mu^{2/3}}$ looks different. There are a few of giants and super-giants in this Table. The values of ratio $\frac{\rho}{\mu^{2/3}}$ are more than 2 for them. It seems that, if to exclude these stars from consideration, the averaging over stars of the main sequence gives value close to 1. Evidently, it needs in more detail consideration.

The luminosity of a star

$$L_0 \sim R_0^2 T_0^4. \quad (128)$$

at taking into account (Eq.121) and (Eq.125) can be expressed as

$$L_0 \sim M^{11/3} \sim M^{3.67} \quad (129)$$

This dependence is shown in Fig.(6) It can be seen that all calculated interdependencies $R(M)$, $T(M)$ and $L(M)$ show a good qualitative agreement with the measuring data. At that it is important, that the quantitative explanation of mass-luminosity dependence discovered at the beginning of 20th century is obtained.




Figure 5: The dependence of the temperature on the star surface over the star mass of close binary stars [8]. Here the temperatures of stars are normalized to surface temperature of the Sun (5875 C), the stars masses are normalized to the mass of Sun. The data are shown on double logarithmic scale. The solid line shows the result of fitting of measurement data ($T_0 \sim M^{0.59}$). The theoretical dependence $T_0 \sim M^{7/12}$ (Eq.125) is shown by dotted line.




Figure 6: The dependence of star luminosity on the star mass of close binary stars [8]. The luminosities are normalized to the luminosity of the Sun, the stars masses are normalized to the mass of the Sun. The data are shown on double logarithmic scale. The solid line shows the result of fitting of measurement data $L \sim M^{3.74}$. The theoretical dependence $L \sim M^{11/3}$ (Eq.129) is shown by dotted line.




Figure 7: The schematic of the star interior

6.3.1 The compilation of the results of calculations

Let us put together the results of calculations. it is energetically favorable for the star to be divided into two areas: the core is located in the central area of the star and the atmosphere is surrounding it from the outside. (Fig.7). The core has the radius:

$$\mathbb{R}_\star = 2.08 \frac{a_B}{Z(A/Z)} \left(\frac{\hbar c}{Gm_p^2} \right)^{1/2} \approx \frac{1.41 \cdot 10^{11}}{Z(A/Z)} cm. \quad (130)$$

It is roughly equal to 1/10 of the stellar radius.

At that the mass of the core is equal to

$$\mathbb{M}_\star = 6.84 \frac{\mathbb{M}_{Ch}}{\left(\frac{A}{Z}\right)^2}. \quad (131)$$

It is almost exactly equal to one half of the full mass of the star.

The plasma inside the core has the constant density

$$n_\star = \frac{16}{9\pi} \frac{Z^3}{a_B^3} \approx 1.2 \cdot 10^{24} Z^3 cm^{-3} \quad (132)$$

and constant temperature

$$\mathbb{T}_\star = \left(\frac{25 \cdot 13}{28\pi^4} \right)^{1/3} \left(\frac{\hbar c}{k a_B} \right) Z \approx Z \cdot 2.13 \cdot 10^7 K. \quad (133)$$

The plasma density and its temperature are decreasing at an approaching to the stellar surface:

$$n_e(r) = n_\star \left(\frac{\mathbb{R}_\star}{r} \right)^6 \quad (134)$$

and

$$T_r = \mathbb{T}_\star \left(\frac{\mathbb{R}_\star}{r} \right)^4. \quad (135)$$

The external radius of the star is determined as

$$\mathbb{R}_0 = \left(\frac{\sqrt{\alpha\pi}}{2\eta} \frac{A}{Z} \frac{m_p}{m_e} \right)^{1/2} \mathbb{R}_\star \approx \frac{6.44 \cdot 10^{11}}{Z(A/Z)^{1/2}} cm \quad (136)$$

and the temperature on the stellar surface is equal to

$$\mathbb{T}_0 = \mathbb{T}_\star \left(\frac{\mathbb{R}_\star}{\mathbb{R}_0} \right)^4 \approx 4.92 \cdot 10^5 \frac{Z}{(A/Z)^2} \quad (137)$$

7 Magnetic fields and magnetic moments of stars

7.1 Magnetic moments of celestial bodies

A thin spherical surface with radius r carrying an electric charge q at the rotation around its axis with frequency Ω obtains the magnetic moment

$$\mathbf{m} = \frac{r^2}{3c} q \Omega. \quad (138)$$

The rotation of a ball charged at density $\varrho(r)$ will induce the magnetic moment

$$\boldsymbol{\mu} = \frac{\Omega}{3c} \int_0^R r^2 \varrho(r) 4\pi r^2 dr. \quad (139)$$

Thus the positively charged core of a star induces the magnetic moment

$$\mathbf{m}_+ = \frac{\sqrt{G} \mathbb{M}_* \mathbb{R}_*^2}{5c} \Omega. \quad (140)$$

A negative charge will be concentrated in the star atmosphere. The absolute value of atmospheric charge is equal to the positive charge of a core. As the atmospheric charge is placed near the surface of a star, its magnetic moment will be more than the core magnetic moment. The calculation shows that as a result, the total magnetic moment of the star will have the same order of magnitude as the core but it will be negative:

$$\mathbf{m}_\Sigma \approx -\frac{\sqrt{G}}{c} \mathbb{M}_* \mathbb{R}_*^2 \Omega. \quad (141)$$

Simultaneously, the torque of a ball with mass $\mathbb{M}$ and radius $\mathbb{R}$ is

$$\mathcal{L} \approx \mathbb{M}_* \mathbb{R}_*^2 \Omega. \quad (142)$$

As a result, for celestial bodies where the force of their gravity induces the electric polarization according to Eq.(40), the giromagnetic ratio will depend on world constants only:

$$\frac{\mathbf{m}_\Sigma}{\mathcal{L}} \approx -\frac{\sqrt{G}}{c}. \quad (143)$$

This relation was obtained for the first time by P.M.S.Blackett [3]. He shows that giromagnetic ratios of the Earth, the Sun and the star 78 Vir are really near to $\sqrt{G}/c$.

By now the magnetic fields, masses, radii and velocities of rotation are known for all planets of the Solar system and for a some stars [15]. These measuring data are shown in Fig.(8), which is taken from [15]. It is possible to see that these data are in satisfactory agreement with Blackett's ratio. At some assumption, the same parameters can be calculated for pulsars. All measured masses of




Figure 8: The observed values of magnetic moments of celestial bodies vs. their angular momenta [15]. In ordinate, the logarithm of the magnetic moment (in $G s \cdot cm^3$) is plotted; in abscissa the logarithm of the angular momentum (in $erg \cdot s$) is shown. The solid line illustrates Eq.(143). The dash-dotted line fits of observed values.

pulsars are equal by the order of magnitude [17]. It is in satisfactory agreement with the condition of equilibrium of relativistic matter (see). It gives a possibility to consider that masses and radii of pulsars are determined. According to generally accepted point of view, pulsar radiation is related with its rotation, and it gives their rotation velocity. These assumptions permit to calculate the giromagnetic ratios for three pulsars with known magnetic fields on their poles [2]. It is possible to see from Fig.(8), the giromagnetic ratios of these pulsars are in agreement with Blackett's ratio.

7.2 Magnetic fields of hot stars

To make the above rough estimation for magnetic fields induced by the star atmosphere more accurate, we can take into account the distribution of electric charges inside the star atmosphere. In accordance with this distribution, the atmosphere at its rotation induces the moment

$$\mathbf{m}_- = -\frac{\sqrt{GM_\star}\Omega\mathbb{R}_\star^2}{9c} \int_{\mathbb{R}_\star}^{R_0} r^2 \frac{d}{dr} \sqrt{\xi^3(2-\xi^3)} dr, \quad (144)$$

Where $\xi = \frac{\mathbb{R}_\star}{r}$.

As $\mathbb{R}_\star \ll R_0$, the magnetic moment of a core can be neglected. In this case, the total magnetic moment of a star

$$\mathbf{m} \approx -2\frac{\sqrt{GM_\star}}{3c} \left(\frac{R_0^{7/2}}{\mathbb{R}_\star^{3/2}} \right) \Omega. \quad (145)$$

and magnetic field near the pole of a star

$$\mathcal{H} \approx -\frac{2\sqrt{GM_\star}}{9c} \frac{R_0^{1/2}}{\mathbb{R}_\star^{3/2}} \Omega. \quad (146)$$

Using of substitution of relations obtained above, one can conclude that the magnetic field on a star pole must not depend on the parameter Z and very slightly depends on A/Z , and this dependence can be neglected. I.e. the calculations show that this field practically is not depending on the mass, on the radius and on the temperature of a hot star and must depend on the velocity of star rotation only:

$$\mathcal{H} \approx -0.47 \left(\frac{m_e}{m_p} \right)^{3/4} \frac{\alpha c}{G^{1/2} \eta^{1/4}} \Omega \approx -8.8 \cdot 10^8 \Omega \quad Oe. \quad (147)$$

The magnetic fields are measured for stars of Ap-class [13]. These stars are characterized by changing their brightness in time. The periods of these changes are measured too. At present there is no full understanding of causes of these visible changes of the luminosity. If these luminosity changes caused by some internal reasons will occur not uniformly on a star surface, one can conclude that the measured period of the luminosity change can depend on star rotation.

It is possible to think that at relatively rapid rotation of a star, the period of a visible change of the luminosity can be determined by this rotation in general. To check this suggestion, we can compare the calculated dependence (Eq.147) with measuring data [13] (see Fig. 9). Evidently one must not expect very good coincidence of calculations and measuring data, because calculations were made for the case of a spherically symmetric model and measuring data are obtained for stars where this symmetry is obviously violated. So getting consent on order of the value can be considered as wholly satisfied. It should be said that Eq.(147) does not working well in case with the Sun. The Sun surface rotates with period $T \approx 25 \div 30$ days. At this velocity of rotation, the magnetic field on the Sun pole calculated accordingly to Eq.(147) must be about 1 kOe. The dipole field of Sun according to experts estimation is approximately 20 times lower. There can be several reasons for that.




Figure 9: The dependence of magnetic fields on poles of Ap-stars as a function of their rotation velocity [13]. The line shows Eq.(147)).

8 The angular velocity of the apsidal rotation in binary stars

8.1 The apsidal rotation of close binary stars

The apsidal rotation (or periastron rotation) of close binary stars is a result of their non-Keplerian movement which originates from the non-spherical form of stars. This non-sphericity has been produced by rotation of stars around their axes or by their mutual tidal effect. The second effect is usually smaller and can be neglected. The first and basic theory of this effect was developed by A.Clairault at the beginning of the XVIII century. Now this effect was measured for approximately 50 double stars. According to Clairault's theory the velocity of periastron rotation must be approximately 100 times faster if matter is uniformly distributed inside a star. Reversely, it would be absent if all star mass is concentrated in the star center. To reach an agreement between the measurement data and calculations, it is necessary to assume that the density of substance grows in direction to the center of a star and here it runs up to a value which is hundreds times greater than mean density of a star. Just the same mass concentration of the stellar substance is supposed by all standard theories of a star interior. It has been usually considered as a proof of astrophysical models. But it can be considered as a qualitative argument. To obtain a quantitative agreement between theory and measurements, it is necessary to fit parameters of the stellar substance distribution in each case separately.

Let us consider this problem with taking into account the gravity induced electric polarization of plasma in a star. As it was shown above, one half of full mass of a star is concentrated in its plasma core at a permanent density. Therefore, the effect of periastron rotation of close binary stars must be reviewed with the account of a change of forms of these star cores.

According to [4],[14] the ratio of the angular velocity ω of rotation of periastron which is produced by the rotation of a star around its axis with the angular velocity Ω is

$$\frac{\omega}{\Omega} = \frac{3}{2} \frac{(I_A - I_C)}{Ma^2} \quad (148)$$

where I_A and I_C are the moments of inertia relatively to principal axes of the ellipsoid. Their difference is

$$I_A - I_C = \frac{M}{5}(a^2 - c^2), \quad (149)$$

where a and c are the equatorial and polar radii of the star.

Thus we have

$$\frac{\omega}{\Omega} \approx \frac{3}{10} \frac{(a^2 - c^2)}{a^2}. \quad (150)$$

8.2 The equilibrium form of the core of a rotating star

In the absence of rotation the equilibrium equation of plasma inside star core (Eq.42 is

$$\gamma \mathbf{g}_G + \rho_G \mathbf{E}_G = 0 \quad (151)$$

where $\gamma, \mathbf{g}_G, \rho_G$ and $\mathbf{E}_G$ are the substance density the acceleration of gravitation, gravity-induced density of charge and intensity of gravity-induced electric field ($div \mathbf{g}_G = 4\pi G \gamma$, $div \mathbf{E}_G = 4\pi \rho_G$ and $\rho_G = \sqrt{G}\gamma$).

One can suppose, that at rotation, under action of a rotational acceleration $\mathbf{g}_\Omega$, an additional electric charge with density ρ_Ω and electric field $\mathbf{E}_\Omega$ can exist, and the equilibrium equation obtains the form:

$$(\gamma_G + \gamma_\Omega)(\mathbf{g}_G + \mathbf{g}_\Omega) = (\rho_G + \rho_\Omega)(\mathbf{E}_G + \mathbf{E}_\Omega), \quad (152)$$

where

$$div (\mathbf{E}_G + \mathbf{E}_\Omega) = 4\pi(\rho_G + \rho_\Omega) \quad (153)$$

or

$$div \mathbf{E}_\Omega = 4\pi\rho_\Omega. \quad (154)$$

We can look for a solution for electric potential in the form

$$\varphi = C_\Omega r^2(3\cos^2\theta - 1) \quad (155)$$

or in Cartesian coordinates

$$\varphi = C_\Omega(3z^2 - x^2 - y^2 - z^2) \quad (156)$$

where C_Ω is a constant.

Thus

$$E_x = 2 C_\Omega x, E_y = 2 C_\Omega y, E_z = -4 C_\Omega z \quad (157)$$

and

$$div \mathbf{E}_\Omega = 0 \quad (158)$$

and we obtain important equations:

$$\rho_\Omega = 0; \quad (159)$$

$$\gamma g_\Omega = \rho \mathbf{E}_\Omega. \quad (160)$$

Since centrifugal force must be contra-balanced by electric force

$$\gamma 2\Omega^2 x = \rho 2C_\Omega x \quad (161)$$

and

$$C_\Omega = \frac{\gamma \Omega^2}{\rho} = \frac{\Omega^2}{\sqrt{G}} \quad (162)$$

The potential of a positive uniform charged ball is

$$\varphi(r) = \frac{Q}{R} \left(\frac{3}{2} - \frac{r^2}{2R^2} \right) \quad (163)$$

The negative charge on the surface of a sphere induces inside the sphere the potential

$$\varphi(R) = -\frac{Q}{R} \quad (164)$$

where according to Eq.(151) $Q = \sqrt{G}M$, and M is the mass of the star.

Thus the total potential inside the considered star is

$$\varphi_\Sigma = \frac{\sqrt{G}M}{2R} \left(1 - \frac{r^2}{R^2} \right) + \frac{\Omega^2}{\sqrt{G}} r^2 (3\cos^2\theta - 1) \quad (165)$$

Since the electric potential must be equal to zero on the surface of the star, at $r = a$ and $r = c$

$$\varphi_\Sigma = 0 \quad (166)$$

and we obtain the equation which describes the equilibrium form of the core of a rotating star (at $\frac{a^2 - c^2}{a^2} \ll 1$)

$$\frac{a^2 - c^2}{a^2} \approx \frac{9}{2\pi} \frac{\Omega^2}{G\gamma}. \quad (167)$$

8.3 The angular velocity of the apsidal rotation

Taking into account of Eq.(167) we have

$$\frac{\omega}{\Omega} \approx \frac{27}{20\pi} \frac{\Omega^2}{G\gamma} \quad (168)$$

If both stars of a close pair induce a rotation of periastron, this equation transforms to

$$\frac{\omega}{\Omega} \approx \frac{27}{20\pi} \frac{\Omega^2}{G} \left(\frac{1}{\gamma_1} + \frac{1}{\gamma_2} \right), \quad (169)$$

where γ_1 and γ_2 are densities of star cores.

The equilibrium density of star cores is known (Eq.(23)):

$$\gamma = \frac{16}{9\pi^2} \frac{A}{Z} m_p \frac{Z^3}{a_B^3}. \quad (170)$$

If we introduce the period of ellipsoidal rotation $P = \frac{2\pi}{\Omega}$ and the period of the rotation of periastron $U = \frac{2\pi}{\omega}$, we obtain from Eq.(168)

$$\frac{P}{U} \left(\frac{P}{T} \right)^2 \approx \sum_1^2 \xi_i, \quad (171)$$

where

$$T = \sqrt{\frac{243 \pi^3}{80}} \tau_0 \approx 10\tau_0, \quad (172)$$

$$\tau_0 = \sqrt{\frac{a_B^3}{G m_p}} \approx 7.7 \cdot 10^2 \text{sec} \quad (173)$$

and

$$\xi_i = \frac{Z_i}{A_i(Z_i + 1)^3}. \quad (174)$$

8.4 The comparison of the calculated angular velocity of the periastron rotation with observations

Because the substance density (Eq.(170)) is depending approximately on the second power of the nuclear charge, the periastron movement of stars consisting of heavy elements will fall out from the observation as it is very slow. Practically the obtained equation (171) shows that it is possible to observe the periastron rotation of a star consisting of light elements only.

The value $\xi = Z/[AZ^3]$ is equal to 1/8 for hydrogen, 0.0625 for deuterium, $1.85 \cdot 10^{-2}$ for helium. The resulting value of the periastron rotation of double stars will be the sum of separate stars rotation. The possible combinations of a couple and their value of $\sum_1^2 \xi_i$ for stars consisting of light elements is shown in Table 8.4.

star1 composed of	star2 composed of	$\xi_1 + \xi_2$
H	H	.25
H	D	0.1875
H	He	0.143
H	hn	0.125
D	D	0.125
D	He	0.0815
D	hn	0.0625
He	He	0.037
He	hn	0.0185

Table 8.4

The "hn" notation in Table 8.4 indicates that the second component of the couple consists of heavy elements or it is a dwarf.

The results of measuring of main parameters for close binary stars are gathered in [8]. For reader convenience, the data of these measurement is applied in




Figure 10: The distribution of close binary stars [8] on value of $(\mathcal{P}/\mathcal{U})(\mathcal{P}/\mathcal{T})^2$. Lines show parameters $\sum_1^2 \xi_i$ for different light atoms in accordance with 174.

the Table in Appendix. One can compare our calculations with data of these measurements. The distribution of close binary stars on value of $(\mathcal{P}/\mathcal{U})(\mathcal{P}/\mathcal{T})^2$ is shown in Fig.10 on logarithmic scale. The lines mark the values of parameters $\sum_1^2 \xi_i$ for different light atoms in accordance with 174. It can be seen that calculated values the periastron rotation for stars composed by light elements which is summarized in Table8.4 are in good agreement with separate peaks of measured data. It confirms that our approach to interpretation of this effect is adequate to produce a satisfactory accuracy of estimations.

9 The solar seismic oscillations

9.1 The spectrum of solar seismic oscillations

The measurements [6] show that the Sun surface is subjected to a seismic vibration. The most intensive oscillations have the period about five minutes and the wave length about 10^4 km or about hundredth part of the Sun radius. Their spectrum obtained by BISON collaboration is shown in Fig.11.

It is supposed, that these oscillations are a superposition of a big number of different modes of resonant acoustic vibrations, and that acoustic waves propagate in different trajectories in the interior of the Sun and they have multiple reflection from surface. With these reflections trajectories of same waves can be closed and as a result standing waves are forming.

Specific features of spherical body oscillations are described by the expansion in series on spherical functions. These oscillations can have a different number of wave lengths on the radius of a sphere (n) and a different number of wave lengths on its surface which is determined by the l -th spherical harmonic. It is accepted to describe the sunny surface oscillation spectrum as the expansion in series [5]:

$$\nu_{nlm} \simeq \Delta\nu_0(n + \frac{l}{2} + \epsilon_0) - l(l+1)D_0 + m\Delta\nu_{rot}. \quad (175)$$

Where the last item is describing the effect of the Sun rotation and is small. The main contribution is given by the first item which creates a large splitting in the spectrum (Fig.11)

$$\Delta\nu = \nu_{n+1,l} - \nu_{n,l}. \quad (176)$$

The small splitting of spectrum (Fig.11) depends on the difference

$$\delta\nu_l = \nu_{n,l} - \nu_{n-1,l+2} \approx (4l+6)D_0. \quad (177)$$

A satisfactory agreement of these estimations and measurement data can be obtained at [5]

$$\Delta\nu_0 = 120 \mu Hz, \epsilon_0 = 1.2, D_0 = 1.5 \mu Hz, \Delta\nu_{rot} = 1 \mu Hz. \quad (178)$$

To obtain these values of parameters $\Delta\nu_0$, ϵ_0 , D_0 from theoretical models is not possible. There are a lot of qualitative and quantitative assumptions used at a model construction and a direct calculation of spectral frequencies transforms into a unresolved complicated problem.

Thus, the current interpretation of the measuring spectrum by the spherical harmonic analysis does not make it clear. It gives no hint to an answer to the question: why oscillations close to hundredth harmonics are really excited and there are no waves near fundamental harmonic?

The measured spectra have a very high resolution (see Fig.(11)). It means that an oscillating system has high quality. At this condition, the system must have oscillation on a fundamental frequency. Some peculiar mechanism must




Figure 11: (a) The power spectrum of solar oscillation obtained by means of Doppler velocity measurement in light integrated over the solar disk. The data were obtained from the BISON network [6]. (b) An expanded view of a part of frequency range.

exist to force a system to oscillate on a high harmonic. The current explanation does not clarify it.

It is important, that now the solar oscillations are measured by means of two different methods. The solar oscillation spectra which was obtained on program "BISON", is shown on Fig.(11)). It has a very high resolution, but (accordingly to the Liouville's theorem) it was obtained with some loss of luminosity, and as a result not all lines are well statistically worked.

Another spectrum was obtained in the program "SOHO/GOLF". Conversely, it is not characterized by high resolution, instead it gives information about general character of the solar oscillation spectrum (Fig.12)).

The existence of this spectrum requires to change the view at all problems of solar oscillations. The theoretical explanation of this spectrum must give answers at least to four questions :

1. Why does the whole spectrum consist from a large number of equidistant spectral lines?
2. Why does the central frequency of this spectrum $\mathcal{F}$ is approximately equal to $\approx 3.23 \text{ mHz}$?
3. Why does this spectrum splitting f is approximately equal to $67.5 \text{ } \mu\text{Hz}$?
4. Why does the intensity of spectral lines decrease from the central line to the periphery?

The answers to these questions can be obtained if we take into account electric polarization of a solar core.

The description of measured spectra by means of spherical analysis does not make clear of the physical meaning of this procedure. The reason of difficulties lies in attempt to consider the oscillations of a Sun as a whole. At existing dividing of a star into core and atmosphere, it is easy to understand that the core oscillation must form a measured spectrum. The fundamental mode of this oscillation must be determined by its spherical mode when the Sun radius oscillates without changing of the spherical form of the core. It gives a most low-lying mode with frequency:

$$\Omega_s \approx \frac{c_s}{\mathbb{R}_\star}, \quad (179)$$

where c_s is sound velocity in the core.

It is not difficult to obtain the numerical estimation of this frequency by order of magnitude. Supposing that the sound velocity in dense matter is 10^7 cm/c and radius is close to $\frac{1}{10}$ of external radius of a star, i.e. about 10^{10} cm , one can obtain as a result

$$F = \frac{\Omega_s}{2\pi} \approx 10^{-3} \text{ Hz} \quad (180)$$

It gives possibility to conclude that this estimation is in agreement with measured frequencies. Let us consider this mechanism in more detail.




Figure 12: (a) The measured power spectrum of solar oscillation. The data were obtained from the SOHO/GOLF measurement [16]. (b) The calculated spectrum described by Eq.(200) at $\langle Z \rangle = 3.4$ and $A/Z = 5$.

9.2 The sound speed in hot plasma

The pressure of high temperature plasma is a sum of the plasma pressure (ideal gas pressure) and the pressure of black radiation:

$$P = n_e kT + \frac{\pi^2}{45\hbar^3 c^3} (kT)^4. \quad (181)$$

and its entropy is

$$S = \frac{1}{\frac{A}{Z} m_p} \ln \frac{(kT)^{3/2}}{n_e} + \frac{4\pi^2}{45\hbar^3 c^3 n_e} (kT)^3, \quad (182)$$

The sound speed c_s can be expressed by Jacobian [9]:

$$c_s^2 = \frac{D(P, S)}{D(\rho, S)} = \frac{\left(\frac{D(P, S)}{D(n_e, T)} \right)}{\left(\frac{D(\rho, S)}{D(n_e, T)} \right)} \quad (183)$$

or

$$c_s = \left\{ \frac{5}{9} \frac{kT}{A/Z m_p} \left[1 + \frac{2 \left(\frac{4\pi^2}{45\hbar^3 c^3} \right)^2 (kT)^6}{5n_e \left[n_e + \frac{8\pi^2}{45\hbar^3 c^3} (kT)^3 \right]} \right] \right\}^{1/2} \quad (184)$$

For $T = T_\star$ and $n_e = n_\star$ we have:

$$\frac{4\pi^2 (kT_\star)^3}{45\hbar^3 c^3 n_\star} \approx 0.18. \quad (185)$$

Finally we obtain:

$$c_s = \left\{ \frac{5}{9} \frac{T_\star}{(A/Z) m_p} [1.01] \right\}^{1/2} \approx 3.14 \cdot 10^7 \left(\frac{Z}{A/Z} \right)^{1/2} \text{ cm/s}. \quad (186)$$

9.3 The basic elastic oscillation of a spherical core

Star cores consist of dense high temperature plasma which is a compressible matter. The basic mode of elastic vibrations of a spherical core is related with its radius oscillation. For the description of this type of oscillation, the potential ϕ of displacement velocities $v_r = \frac{\partial \psi}{\partial r}$ can be introduced and the motion equation can be reduced to the wave equation expressed through ϕ [9]:

$$c_s^2 \Delta \phi = \ddot{\phi}, \quad (187)$$

and a spherical derivative for periodical in time oscillations ($\sim e^{-i\Omega_s t}$) is:

$$\Delta \phi = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \phi}{\partial r} \right) = -\frac{\Omega_s^2}{c_s^2} \phi. \quad (188)$$

It has the finite solution for the full core volume including its center

$$\phi = \frac{A}{r} \sin \frac{\Omega_s r}{c_s}, \quad (189)$$

where A is a constant. For small oscillations, when displacements on the surface u_R are small ($u_R/R = v_R/\Omega_s R \rightarrow 0$) we obtain the equation:

$$\text{tg} \frac{\Omega_s \mathbb{R}}{c_s} = \frac{\Omega_s \mathbb{R}}{c_s} \quad (190)$$

which has the solution:

$$\frac{\Omega_s \mathbb{R}}{c_s} \approx 4.49. \quad (191)$$

Taking into account Eq.(186)), the main frequency of the core radial elastic oscillation is

$$\Omega_s = 4.49 \left\{ 1.4 \left[\frac{Gm_p}{r_B^3} \right] \frac{A}{Z} \left(Z + 1 \right)^3 \right\}^{1/2}. \quad (192)$$

It can be seen that this frequency depends on Z and A/Z only. Some values of frequencies of radial sound oscillations $\mathcal{F} = \Omega_s/2\pi$ calculated from this equation for selected A/Z and Z are shown in third column of Table (9.3).

Table (9.3)

Z	A/Z	$\mathcal{F}, mHz$ (calculated (192))	star	$\mathcal{F}, mHz$ measured
1	1	0.23	$\xi Hydrae$	~ 0.1
1	2	0.32	$\nu Indus$	0.3
2	2	0.9	$\eta Bootis$	0.85
2	3	1.12	The Procion($A\alpha CMi$)	1.04
			$\beta Hydrae$	1.08
3	4	2.38	$\alpha Cen A$	2.37
3	5	2.66		
3.4	5	3.24	The Sun	3.23
4	5	4.1		

The star mass spectrum (.1) shows that the ratio A/Z must be ≈ 5 for the Sun. It is in accordance with the calculated frequency of solar core oscillations if the averaged charge of nuclei $Z \approx 3.4$. It is not a confusing conclusion, because the plasma electron gas prevents the decay of β -active nuclei (see Sec.10). This mechanism can probably to stabilize neutron-excess nuclei.

9.4 The low frequency oscillation of the density of a neutral plasma

Hot plasma has the density n_* at its equilibrium state. The local deviations from this state induce processes of density oscillation since plasma tends to return to its steady-state density. If we consider small periodic oscillations of core radius

$$R = \mathbb{R} + u_R \cdot \sin \omega_{n_*} t, \quad (193)$$

where a radial displacement of plasma particles is small ($u_R \ll \mathbb{R}$), the oscillation process of plasma density can be described by the equation

$$\frac{d\mathcal{E}}{dR} = \mathbb{M} \ddot{R}. \quad (194)$$

Taking into account

$$\frac{d\mathcal{E}}{dR} = \frac{d\mathcal{E}_{plasma}}{dn_e} \frac{dn_e}{dR} \quad (195)$$

and

$$\frac{3}{8} \pi^{3/2} \mathbb{N}_e \frac{e^3 a_0^{3/2}}{(k\mathbb{T})^{1/2}} \frac{n_*}{\mathbb{R}^2} = \mathbb{M} \omega_{n_*}^2 \quad (196)$$

From this we obtain

$$\omega_{n_*}^2 = \frac{3}{\pi^{1/2}} k\mathbb{T} \left(\frac{e^2}{a_B k\mathbb{T}} \right)^{3/2} \frac{Z^3}{\mathbb{R}^2 A / Z m_p} \quad (197)$$

and finally

$$\omega_{n_*} = \left\{ \frac{2^8}{3^5} \frac{\pi^{1/2}}{10^{1/2}} \alpha^{3/2} \left[\frac{Gm_p}{a_B^3} \right] \frac{A}{Z} Z^{4.5} \right\}^{1/2}, \quad (198)$$

where $\alpha = \frac{e^2}{\hbar c}$ is the fine structure constant. These low frequency oscillations of neutral plasma density are similar to phonons in solid bodies. At that oscillations with multiple frequencies $k\omega_{n_*}$ can exist. Their power is proportional to $1/\kappa$, as the occupancy these levels in energy spectrum must be reversely proportional to their energy $k\hbar\omega_{n_*}$. As result, low frequency oscillations of plasma density constitute set of vibrations

$$\sum_{\kappa=1} \frac{1}{\kappa} \sin(\kappa\omega_{n_*}t). \quad (199)$$

9.5 The spectrum of solar core oscillations

The set of the low frequency oscillations with ω_η can be induced by sound oscillations with Ω_s . At that, displacements obtain the spectrum:

$$u_R \sim \sin \Omega_s t \cdot \sum_{\kappa=0} \frac{1}{\kappa} \sin \kappa\omega_{n_*} t \sim \xi \sin \Omega_s t + \sum_{\kappa=1} \frac{1}{\kappa} \sin (\Omega_s \pm \kappa\omega_{n_*})t, \quad (200)$$

where ξ is a coefficient ≈ 1 .

This spectrum is shown in Fig.(12).

The central frequency of experimentally measured distribution of solar oscillations is approximately equal to (Fig.(11))

$$\mathcal{F}_\odot \approx 3.23 \text{ mHz} \quad (201)$$

and the experimentally measured frequency splitting in this spectrum is approximately equal to

$$f_\odot \approx 68 \text{ } \mu\text{Hz}. \quad (202)$$

A good agreement of the calculated frequencies of basic modes of oscillations (from Eq.(192) and Eq.(151)) with measurement can be obtained at $Z = 3.4$ and $A/Z = 5$:

$$\mathcal{F}_{Z=3.4; \frac{A}{Z}=5} = \frac{\Omega_s}{2\pi} = 3.24 \text{ mHz}; \quad f_{Z=3.4; \frac{A}{Z}=5} = \frac{\omega_{n_*}}{2\pi} = 68.1 \text{ } \mu\text{Hz}. \quad (203)$$

10 Addendum I: A mechanism of stabilization for neutron-excess nuclei in plasma

10.1 Neutron-excess nuclei and the neutronization

The form of the star mass spectrum (Fig.1) indicates that plasma of many stars consists of neutron-excess nuclei with the ratio $A/Z = 3, 4, 5$ and so on. These nuclei are subjects of a decay under the "terrestrial" conditions. Hydrogen isotopes ${}^4_1H$, ${}^5_1H$, ${}^6_1H$ have short time of life and emit electrons with energy more than 20 Mev. The decay of helium isotopes ${}^6_2He$, ${}^8_2He$, ${}^{10}_2He$ have the times of life, which can reach tenth part of the seconds.

Stars have the time of life about billions years and the lines of their spectrum of masses are not smoothed. Thus we should suppose that there is some mechanism of stabilization of neutron-excess nuclei inside stars. This mechanism is well known - it is neutronization [9]§106. It is accepted to think that this mechanism is characteristic for dwarfs with density of particles about 10^{30} per cm^3 and pressure of relativistic electron gas

$$P \approx \hbar c \cdot n_e^{4/3} \approx 10^{23} \text{ dyne/cm}^2. \quad (204)$$

The possibility of realization of neutronization in dense plasma is considered below in detail. At thus, we must try to find an explanation to characteristic features of the star mass spectrum. At first, we can see that, there is actually quite a small number of stars with $A/Z = 2$ exactly. The question is arising: why there are so few stars, which are composed by very stable nuclei of helium-4? At the same time, there are many stars with $A/Z = 4$, i.e. consisting apparently of a hydrogen-4, as well as stars with $A/Z = 3/2$, which hypothetically could be composed by another isotope of helium - helium-3.

10.1.1 The electron cloud of plasma cell

Let us consider a possible mechanism of the action of the electron gas effect on the plasma nuclear subsystem. It is accepted to consider dense plasma to be divided in plasma cells. These cells are filled by electron gas and they have positively charged nuclei in their centers [12].

This construction is non stable from the point of view of the classical mechanics because the opposite charges collapse is "thermodynamic favorable". To avoid a divergence in the theoretical description of this problem, one can artificially cut off the integrating at some small distance characterizing the particles interaction. For example, nuclei can be considered as hard cores with the finite radii.

It is more correctly, to consider this structure as a quantum-mechanical object and to suppose that the electron can not approach the nucleus closer than its own de Broglie's radius λ_e .

Let us consider the behavior of the electron gas inside the plasma cell. If to express the number of electrons in the volume V through the density of electron

n_e , then the maximum value of electron momentum [9]:

$$p_F = (3\pi^2 n_e)^{1/3} \hbar. \quad (205)$$

The kinetic energy of the electron gas can be founded from the general expression for the energy of the Fermi-particles, which fills the volume V [9]:

$$\mathcal{E} = \frac{Vc}{\pi^2 \hbar^3} \int_0^{p_F} p^2 \sqrt{m_e^2 c^2 + p^2} dp. \quad (206)$$

After the integrating of this expression and the subtracting of the energy at rest, we can calculate the kinetic energy of the electron:

$$\mathcal{E}_{kin} = \frac{3}{8} m_e c^2 \left[\frac{\xi(2\xi^2 + 1)\sqrt{\xi^2 + 1} - \text{Arcsinh}(\xi) - \frac{8}{3}\xi^3}{\xi^3} \right] \quad (207)$$

(where $\xi = \frac{p_F}{m_e c}$).

The potential energy of an electron is determined by the value of the attached electric field. The electrostatic potential of this field $\varphi(r)$ must be equal to zero at infinity.⁷ With this in mind, we can write the energy balance equation of electron

$$\mathcal{E}_{kin} = e\varphi(r). \quad (208)$$

The potential energy of an electron at its moving in an electric field of the nucleus can be evaluated on the basis of the Lorentz transformation [11]§24. If in the laboratory frame of reference, where an electric charge placed, it creates an electric potential φ_0 , the potential in the frame of reference moving with velocity v is

$$\varphi = \frac{\varphi_0}{\sqrt{1 - \frac{v^2}{c^2}}}. \quad (209)$$

Therefore, the potential energy of the electron in the field of the nucleus can be written as:

$$\mathcal{E}_{pot} = -\frac{Ze^2}{r} \frac{\xi}{\beta}. \quad (210)$$

Where

$$\beta = \frac{v}{c}. \quad (211)$$

and

$$\xi \equiv \frac{p}{m_e c}, \quad (212)$$

m_e is the mass of electron in the rest.

⁷In general, if there is an uncompensated electric charge inside the cell, then we would have to include it to the potential $\varphi(r)$. However, we can do not it, because will consider only electro-neutral cell, in which the charge of the nucleus exactly offset by the electronic charge, so the electric field on the cell border is equal to zero.

And one can rewrite the energy balance Eq.(208) as follows:

$$\frac{3}{8}m_e c^2 \xi \mathbb{Y} = e\varphi(r) \frac{\xi}{\beta}. \quad (213)$$

where

$$\mathbb{Y} = \left[\frac{\xi(2\xi^2 + 1)\sqrt{\xi^2 + 1} - \text{Arcsinh}(\xi) - \frac{8}{3}\xi^3}{\xi^4} \right]. \quad (214)$$

Hence

$$\varphi(r) = \frac{3}{8} \frac{m_e c^2}{e} \beta \mathbb{Y}. \quad (215)$$

In according with Poisson's electrostatic equation

$$\Delta\varphi(r) = 4\pi en_e \quad (216)$$

or at taking into account that the electron density is depending on momentum (Eq.(205)), we obtain

$$\Delta\varphi(r) = \frac{4e}{3\pi} \left(\frac{\xi}{\tilde{\chi}_C} \right)^3, \quad (217)$$

where $\tilde{\chi}_C = \frac{\hbar}{m_e c}$ is the Compton radius.

At introducing of the new variable

$$\varphi(r) = \frac{\chi(r)}{r}, \quad (218)$$

we can transform the Laplacian:

$$\Delta\varphi(r) = \frac{1}{r} \frac{d^2\chi(r)}{dr^2}. \quad (219)$$

As (Eq.(215))

$$\chi(r) = \frac{3}{8} \frac{m_e c^2}{e} \mathbb{Y} \beta r, \quad (220)$$

the differential equation can be rewritten:

$$\frac{d^2\chi(r)}{dr^2} = \frac{\chi(r)}{\mathbb{L}^2}, \quad (221)$$

where

$$\mathbb{L} = \left(\frac{9\pi}{32} \frac{\mathbb{Y}\beta}{\alpha\xi^3} \right)^{1/2} \tilde{\chi}_C, \quad (222)$$

$\alpha = \frac{1}{137}$ is the fine structure constant.

This differential equation has the solution:

$$\chi(r) = C \cdot \exp\left(-\frac{r}{\mathbb{L}}\right). \quad (223)$$

Thus, the equation of equilibrium of the electron gas inside a cell (Eq.(213)) obtains the form:

$$\frac{Ze}{r} \cdot e^{-r/\mathbb{L}} = \frac{3}{8} m_e c^2 \beta \mathbb{Y}. \quad (224)$$

10.2 The Thomas-Fermi screening

Let us consider the case when an ion is placed at the center of a cell, the external shells don't permit the plasma electron to approach to the nucleus on the distances much smaller than the Bohr radius. The electron moving is non-relativistic in this case. At that $\xi \rightarrow 0$, the kinetic energy of the electron

$$\mathcal{E}_{kin} = \frac{3}{8}m_e c^2 \xi \mathbb{Y} \rightarrow \frac{3}{5}E_F \quad , \quad (225)$$

and the screening length

$$\mathbb{L} \rightarrow \sqrt{\frac{\mathcal{E}_F}{6\pi e^2 n_e}}. \quad (226)$$

Thus, we get the Thomas-Fermi screening in the case of the non-relativistic motion of an electron.

10.3 The screening with relativistic electrons

In the case the «bare» nucleus, there is nothing to prevent the electron to approach it at an extremely small distance λ_{min} , which is limited by its own than its de Broglie's wavelength. Its movement in this case becomes relativistic at $\beta \rightarrow 1$ $\xi \gg 1$. In this case, at not too small ξ , we obtain

$$\mathbb{Y} \approx 2 \left(1 - \frac{4}{3\xi} \right), \quad (227)$$

and at $\xi \gg 1$

$$\mathbb{Y} \rightarrow 2. \quad (228)$$

In connection with it, at the distance $r \rightarrow \lambda_{min}$ from a nucleus, the equilibrium equation (224) reforms to

$$\lambda_{min} \simeq Z\alpha\lambda_C \quad . \quad (229)$$

and the density of electron gas in a layer of thickness λ_{min} can be determined from the condition of normalization. As there are Z electrons into each cell, so

$$Z \simeq n_e^\lambda \cdot \lambda_{min}^3 \quad (230)$$

From this condition it follows that

$$\xi_\lambda \simeq \frac{1}{2\alpha Z^{2/3}} \quad (231)$$

Where n_e^λ and ξ_λ are the density of electron gas and the relative momentum of electrons at the distance λ_{min} from the nucleus. In accordance with Eq.(207), the energy of all the Z electrons in the plasma cell is

$$\mathcal{E} \simeq Zm_e c^2 \xi_\lambda \quad (232)$$

At substituting of Eq.(231), finally we obtain the energy of the electron gas in a plasma cell:

$$\mathcal{E} \simeq \frac{m_e c^2}{2\alpha} Z^{1/3} \quad (233)$$

This layer provides the pressure on the nucleus:

$$P \simeq \mathcal{E} \left(\frac{\xi}{\lambda_C} \right)^3 \approx 10^{23} \text{dyne/cm}^2 \quad (234)$$

This pressure is in order of value with pressure of neutronization (204).

10.4 The neutronization

. The considered above $\ll$ attachment $\gg$ of the electron to the nucleus in a dense plasma should lead to a phenomenon neutronization of the nucleus, if it is energetically favorable. The $\ll$ attached $\gg$ electron layer should have a stabilizing effect on the neutron-excess nuclei, i.e. the neutron-excess nucleus, which is instable into substance with the atomic structure, will became stable inside the dense plasma. It explains the stable existence of stars with a large ratio of A/Z .

These formulas allow to answer questions related to the characteristics of the star mass distribution. The numerical evaluation of energy the electron gas in a plasma cell gives:

$$\mathcal{E} \simeq \frac{m_e c^2}{2\alpha} Z^{1/3} \approx 5 \cdot 10^{-5} Z^{1/3} \text{erg} \quad (235)$$

The mass of nucleus of helium-4 $M({}_2^4\text{He}) = 4.0026 \text{ a.e.m.}$, and the mass hydrogen-4 $M({}_1^4\text{H}) = 4.0278 \text{ a.e.m.}$. The mass defect $\approx 3.8 \cdot 10^{-5} \text{egr}$. Therefore, the reaction



is energetically favorable. At this reaction the nucleus captures the electron from gas and proton becomes a neutron.

There is the visible line of stars with the ratio $A/Z = 3/2$ at the star mass spectrum. It can be attributed to the stars, consisting of ${}_2^3\text{He}$, ${}_4^6\text{Be}$, ${}_6^9\text{C}$, etc.

It is not difficult to verify at direct calculation that the reactions of neutronization and transforming of ${}_2^3\text{He}$ into ${}_1^3\text{H}$ and ${}_4^6\text{Be}$ into ${}_3^6\text{Li}$ are energetically allowed. So the nuclei ${}_2^3\text{He}$ and ${}_4^6\text{Be}$ should be conversed by neutronization into ${}_1^3\text{H}$ and ${}_3^6\text{Li}$. The line $A/Z = 3/2$ of the star mass spectrum can not be formed by these nuclei. However, the reaction



is not energetically allowed and therefore it is possible to believe that the stars of the above line mass spectrum may consist of carbon-9.

It is interesting, is it possible to study this effect under laboratory conditions? One can use gaseous tritium for it and can measure the rate of its decay at two states. First the rate of tritium decay at atomic (molecular) state. Second, the same rate under the effect of ionizing electric discharge. Essentially, one must aim to use the dense gas and the power discharge in order to ionize more atoms of tritium. Taking into account the preceding consideration, such a test does not look as a hopeless project.

11 Addendum II: Other stars, their classification and some cosmology

The Schwarzsprung-Rassel diagram is now a generally accepted base for star classification. It seems that a classification based on the EOS of substance may be more justified from physical point of view. It can be emphasized by possibility to determine the number of classes of celestial bodies.

The matter can totally have eight states (Fig.(13)).

The atomic substance at low temperature exists as condensed matter (solid or liquid). At high temperature it transforms into gas phase.

The electron -nuclear plasma can exist in four states. It can be relativistic or non-relativistic. The electron gas of non-relativistic plasma can be degenerate (cold) or non-degenerate (hot). The relativistic electron gas is degenerate at temperature below T_F . Very high temperature can remove the degeneration even for relativistic electrons (if, certainly, electronic gas was not ultra-relativistic originally).

In addition to that, a substance can exist as neutron matter with the nuclear density approximately.

At present, assumptions about existence of matter at different states, other than the above-named, seem unfounded. Thus, these above-named states of matter show a possibility of classification of celestial bodies in accordance with this dividing.

11.1 The atomic substance

11.1.1 Small bodies

Small celestial bodies - asteroids and satellites of planets - are usually considered as bodies consisting from atomic matter.

11.1.2 Giants

The transformation of atomic matter into plasma can be induced by action of high pressure, high temperature or both these factors. If these factors inside a body are not high enough, atomic substance can transform into gas state. The characteristic property of this celestial body is absence of electric polarization inside it. If temperature of a body is below ionization temperature of atomic substance but higher than its evaporation temperature, the equilibrium equation comes to

$$-\frac{dP}{dr} = \frac{G\gamma}{r^2}M_r \approx \frac{P}{R} \approx \frac{\gamma}{m_p} \frac{kT}{R}. \quad (238)$$

Thus, the radius of the body

$$R \approx \frac{GMm_p}{kT}. \quad (239)$$

	Low T	High T
Atomic substance	Solid body	gas
Plasma $p_F \ll m_e c$	Non-relativistic degenerate	Non-relativistic Non-degenerate
Plasma $p_F \approx m_e c$	relativistic degenerate	relativistic non-degenerate
neutron matter $p_F \approx m_n c$	relativistic degenerate	relativistic non-degenerate

Figure 13: The steady states of substance

If its mass $M \approx 10^{33} g$ and temperature at its center $T \approx 10^5 K$, its radius is $R \approx 10^2 R_\odot$. These properties are characteristic for giants, where pressure at center is about $P \approx 10^{10} \text{ din/cm}^2$ and it is not enough for substance ionization.

11.2 Plasmas

11.2.1 The non-relativistic non-degenerate plasma. Stars.

Characteristic properties of hot stars consisting of non-relativistic non-degenerate plasma was considered above in detail. Its EOS is ideal gas equation.

11.2.2 Non-relativistic degenerate plasma. Planets.

At cores of large planets, pressures are large enough to transform their substance into plasma. As temperatures are not very high here, it can be supposed, that this plasma can degenerate:

$$T \ll T_F \quad (240)$$

The pressure which is induced by gravitation must be balanced by pressure of non-relativistic degenerate electron gas

$$\frac{GM^2}{6RV} \approx \frac{(3\pi^2)^{2/3}}{5} \frac{\hbar^2}{m_e} \left(\frac{\gamma}{m_p A/Z} \right)^{5/3} \quad (241)$$

It opens a way to estimate the mass of this body:

$$M \approx M_{Ch} \left(\frac{\hbar}{mc} \right)^{3/2} \left(\frac{\gamma}{m_p} \right)^{1/2} \frac{6^{3/2} 9\pi}{4(A/Z)^{5/2}} \quad (242)$$

At density about $\gamma \approx 1 g/cm^3$, which is characteristic for large planets, we obtain their masses

$$M \approx 10^{-3} \frac{M_{Ch}}{(A/Z)^{5/2}} \approx \frac{4 \cdot 10^{30}}{(A/Z)^{5/2}} g \quad (243)$$

Thus, if we suppose that large planets consist of hydrogen ($A/Z=1$), their masses must not be over $4 \cdot 10^{30} g$. It is in agreement with the Jupiter's mass, the biggest planet of the Sun system.

11.2.3 The cold relativistic substance

The kinetic energy of N relativistic particles (207):

$$\mathcal{E}^{kinetic} = \frac{3}{8} N m c^2 \left[\frac{\xi(2\xi^2 + 1) \sqrt{\xi^2 + 1} - \text{Arcsinh}(\xi) - \frac{8}{3}\xi^3}{\xi^3} \right] \quad (244)$$

where $\xi = \frac{p_F}{mc}$ and density of substance:

$$n = \frac{p_F^3}{3\pi^2 \hbar^3} = \frac{\xi^3}{3\pi^2} \left(\frac{mc}{\hbar} \right)^3. \quad (245)$$

The pressure of this system is

$$P = - \left(\frac{d\mathcal{E}^{kinetic}}{dV} \right)_{S=0} = \frac{mc^2}{8\pi^2} \left(\frac{mc}{\hbar} \right)^3 \left[\xi \left(\frac{2}{3}\xi^2 - 1 \right) \sqrt{\xi^2 + 1} + \text{Arcsinh}(\xi) \right] \quad (246)$$

For simplicity we can suppose that main part of mass of relativistic star is concentrated into its core and it is uniformly distributed here. In this case we can write the pressure balance equation:

$$\frac{GM^2}{2R} = 3PV. \quad (247)$$

It permits to estimate the full number of particles into the considered system:

$$N = \left(\frac{\hbar c}{Gm_p^2} \right)^{3/2} \left(\frac{3}{4\pi} \right)^2 \frac{1}{\pi^{3/2}} \left[\frac{\xi \left(\frac{2}{3}\xi^2 - 1 \right) \sqrt{\xi^2 + 1} + \text{Arcsinh}(\xi)}{\xi^4} \right]^{3/2}. \quad (248)$$

Accordingly to the virial theorem:

$$\mathcal{E}_{total} = -\mathcal{E}^{kinetic} \quad (249)$$

and (244), the full energy of a relativistic star

$$\begin{aligned} \mathcal{E}_{total} \sim & -\frac{1}{\xi^9} \left[\xi(2\xi^2 + 1)\sqrt{\xi^2 + 1} - \text{Arcsinh}(\xi) - \frac{8}{3}\xi^3 \right] \cdot \\ & \cdot \left[\xi \left(\frac{2}{3}\xi^2 - 1 \right) \sqrt{\xi^2 + 1} + \text{Arcsinh}(\xi) \right]^{3/2}. \end{aligned} \quad (250)$$

Because this expression has a minimum at $\xi \approx 1.35$, a star consisting of relativistic substance must have the equilibrium particle density:

$$n_\star \approx 8.3 \cdot 10^{-2} \left(\frac{mc}{\hbar} \right)^3. \quad (251)$$

The relativistic degenerate plasma. Dwarfs.

The degenerate relativistic plasma has a cold relativistic electron subsystem. Its nuclear subsystem can be non-relativistic. The star consisting of this plasma accordingly to Eq.(251) at $m = m_e$, must have an energy minimum at electron density

$$n_r \approx 1.4 \cdot 10^{30} \text{cm}^{-3} \quad (252)$$

at a star radius

$$R \approx 10^{-2} R_\odot \quad (253)$$

It is not difficult to see that these densities and radii are characteristic for dwarfs.

The neutron matter. Pulsars. Dwarfs may be considered as stars where a process of neutronization is just beginning. At a nuclear density, plasma turns into neutron matter.⁸

⁸At nuclear density neutrons and protons are indistinguishable inside pulsars as inside a huge nucleus. It permits to suppose a possibility of gravity induced electric polarization in this matter.




Figure 14: The mass distribution of pulsars from binary systems [17]. On abscissa the logarithm of pulsar mass in solar mass is shown.

In accordance with Eq.(251) at $m = m_n$, a star consisting of neutron matter must have the minimal energy at neutron density $n_\star \approx 9 \cdot 10^{39} \text{ particle/cm}^3$. As result the masses of neutron stars must be approximately equal to M_{Ch} . The measured mass distribution of pulsar composing binary stars [17] is shown on Fig.14. It can be considered as a confirmation of the last conclusion.

11.2.4 The hot relativistic plasma.Quasars

Plasma is hot if its temperature is higher than degeneration temperature of its electron gas. The ratio of plasma temperature in the core of a star to the temperature of degradation of its electron gas for case of non-relativistic hot star plasma is (Eq.(28))

$$\frac{T_\star}{T_F(n_\star)} \approx 40 \quad (254)$$

it can be supposed that the same ratio must be characteristic for the case of a relativistic hot star. At this temperature, the radiation pressure plays a main




Figure 15: The mass distribution of galaxies. [1]. On the abscissa, the logarithm of the galaxy mass over the Sun mass is shown.

role and accordingly the equation of the pressure balance takes the form:

$$\frac{GM^2}{6RV} \approx \frac{\pi^2 (kT_\star)^4}{45 (\hbar c)^3} \approx \left(\frac{T_\star}{T_F}\right)^3 kTn \quad (255)$$

This makes it possible to estimate the mass of a hot relativistic star

$$M \approx \left(\frac{T_\star}{T_F}\right)^6 \left(\frac{\hbar c}{Gm_p^2}\right)^{3/2} m_p \approx 10^9 M_\odot \quad (256)$$

According to the existing knowledge, among compact celestial objects only quasars have masses of this level. Apparently it is an agreed-upon opinion that quasars represent some relatively short stage of evolution of galaxies. If we adhere to this hypothesis, the lack of information about quasar mass distribution can be replaced by the distribution of masses of galaxies [1](Fig.15). It can be seen, that this distribution is in a qualitative agreement with supposition that quasars are composed from the relativistic hot plasma.

Certainly the used estimation Eq.(254) is very arbitrary. It is possible to expect the existing of quasars which can have substantially lesser masses.

As the steady-state particle density of relativistic matter n_r is known (Eq.(252)), one can estimate the quasar radius:

$$R_{qu} \approx \sqrt[3]{\frac{M_{qu}}{n_r m_p}} \approx 10^{12} \text{cm}. \quad (257)$$

It is in agreement with the astronomer measuring data obtained from periods of the their luminosity changing.

11.2.5 About the cosmic object categorizations

Thus, it seems possible under some assumptions to find characteristic parameters of different classes of stars, if to proceed from EOS of atomic, plasma and neutron substances. The EOS types can be compared with classes of celestial bodies. As any other EOS are unknown, it gives a reason to think that all classes of celestial bodies are discovered (Fig.16).

11.3 Several words about star evolution

There are not formulas in this section, which could serve as a handhold for suggestions. The formulas of the previous sections can not help too in understanding that, as the evolution of the stars can proceed and as their transitions them from the one class to another are realized, since these formulas were received for the description of the star steady-state. There the comparison of the schemes to the substance categorization (Fig.(13)) and the categorization of star classes (Fig.(16)) can serve as the base of next suggestions.

At analyzing of these schemes, one can suppose that the evolution of stellar objects is going at the reduction of their temperature. In light of it, it is possible to suppose the existing of one more body, from which the development began. Really, neutron matter at nuclear density (Eq.(251)) is not ultra-relativistic. At this density, corresponding to $p_F \approx mc$, its energy and pressure can depend on the temperature if this temperature it is high enough.⁹ It seems, there is no thermodynamical prohibition to imagine this matter so hot when the neutron gas is non-degenerate. An estimation shows that it can be possible if mass of this body is near to $10^{50}g$ or even to $10^{55}g$. As it is accepted to think that full mass of the Universe is about $10^{53}g$, it can be assumed that on an early stage of development of Universe, there was some celestial body with the mass about $10^{53}g$ composed by the neutron matter at the approximately nuclear density with at the temperature above $10^{12}K$. After some time, with temperature decreased it has lost its stability and decayed into quasars with mass up to $10^{12}M_{Ch}$, consisting of the non-degenerate relativistic plasma at $T > 10^{10}K$. After next cooling at loosing of stability they was decaying on galaxies of hot stars with mass about $M \approx M_{Ch}$ and core temperature about $T \approx 10^7K$, composed by non-relativistic hot plasma. A next cooling must leads hot stars to decaying on dwarfs, pulsars, planets or may be on small bodies. The substances of these bodies (in their cores) consists of degenerate plasma (degenerate electron subsystem and cold nuclear subsystem) or cold neutron matter, it makes them stable in expanding and cooling Universe.¹⁰ It is

⁹The ultra-relativistic matter with $p_F \gg mc$ is possessed by limiting pressure which is not depending on temperature.

¹⁰The temperature of plasma inside these bodies can be really quite high as electron gas into dwarfs, for example, will be degenerate even at temperature $T \approx 10^9K$.

	Low T	High T
Atomic substance	asteroid	gaint
Plasma $p_F \ll m_e c$	planet	star
Plasma $p_F \approx m_e c$	dwarf	quasar
neutron matter $p_F \approx m_n c$	pulsar	?

Figure 16: The categorization of celestial bodies

important to emphasize, that the gravity induced electric polarization excludes the possibility of the gravitational collapse as the last stage of the star evolutions.

11.4 About «black holes»

It seems that the idea about the «black holes» existence is organic related to the suggestion about an inevitable collapse of large cosmic bodies on the last stage of their evolutions. However, the models of collapsing masses were appearing as a consequence of the rejection from attention of the gravity induced electric polarization of the intra-stellar plasma, which was considered in previous chapters. If to take into account this mechanism, the possibility of collapse must be excluded. It allows newly to take a look on the «black hole» problem.

In accordance with the standard approach, the Schwarzschild radius of «black hole» with M is

$$r_{bh} = \frac{2GM}{c^2} \quad (258)$$

and accordingly the average density of «black holes»:

$$\gamma_{bh} = \frac{3c^6}{32\pi M^2 G^3}. \quad (259)$$

The estimations which was made in previous chapters are showing that all large inwardly-galactic objects of all classes - a stars, dwarves, pulsars, giants - possess the mass of the order M_{Ch} (or $10M_{Ch}$). As the density of these objects are small relatively to the limit (259). As result, a searching of «black hole» inside stellar objects of our Galaxy seems as hopeless.

On the other hand, stellar objects, consisting of hot relativistic plasma - a quasars, in accordance with their mass and density, may stay «black holes». The process of collapse is not needed for their creation. As the quasar mass $M_{qu} \gg M_{Ch}$, all other stellar objects must organize their moving around it and one can suppose that a «black hole» can exist at the center of our Galaxy.

12 The conclusion

Evidently, the main conclusion from the above consideration consists in statement of the fact that now there are quite enough measuring data to place the theoretical astrophysics on a reliable foundation. All above measuring data are known for a relatively long time. The traditional system of view based on the Euler equation in the form (1) could not give a possibility to explain and even to consider, with due proper attention, to these data. Taking into account the gravity induced electric polarization of plasma and a change starting postulate gives a possibility to obtain results for explanation of measuring data considered above.

Basically these results are the following.

Using the standard method of plasma description leads to the conclusion that at conditions characteristic for the central stellar region, the plasma has the minimum energy at constant density n_* (Eq.(23)) and at the constant temperature T_* (Eq.(79)).

This plasma forms the core of a star, where the pressure is constant and gravity action is balanced by the force of the gravity induced by the electric polarization. The virial theorem gives a possibility to calculate the stellar core mass M_* (Eq.(83)) and its radius R_* (85). At that the stellar core volume is approximately equal to 1/1000 part of full volume of a star.

The remaining mass of a star located over the core has a density approximately thousand times smaller and it is convenient to name it a star atmosphere. At using thermodynamical arguments, it is possible to obtain the radial dependence of plasma density inside the atmosphere $n_a \approx r^{-6}$ (Eq.(55)) and the radial dependence of its temperature $T_a \approx r^{-4}$ (Eq.(56)).

It gives a possibility to conclude that the mass of the stellar atmosphere M_a (Eq.(57)) is almost exactly equal to the stellar core mass. Thus, the full stellar mass can be calculated. It depends on the ratio of the mass and the charge of nuclei composing the plasma. This claim is in a good agreement with the measuring data of the mass distribution of both - binary stars and close binary stars (Fig.(1)-(2)) ¹¹. At that it is important that the upper limit of masses of both - binary stars and close binary stars - is in accordance with the calculated value of the mass of the hydrogen star (Eq.(84)). The obtained formula explains the origin of sharp peaks of stellar mass distribution - they evidence that the substance of these stars have a certain value of the ratio A/Z . In particular the solar plasma according to (Eq.(84)) consists of nuclei with $A/Z = 5$.

¹¹The measurement of parameters of these stars has a satisfactory accuracy only.

Knowing temperature and substance density on the core and knowing their radial dependencies, it is possible to estimate the surface temperature T_0 (96) and the radius of a star R_0 (95). It turns out that these measured parameters must be related to the star mass with the ratio $T_0 R_0 \sim M^{5/4}$ (104). It is in a good agreement with measuring data (Fig.(3)).

Using another thermodynamical relation - the Poisson's adiabat - gives a way to determine the relation between radii of stars and their masses $R_0^3 \sim M^2$ (Eq.(121)), and between their surface temperatures and masses $T_0 \sim M^{5/7}$ (Eq.(125)). It gives the quantitative explanation of the mass-luminosity dependence (Fig.(6)).

According to another familiar Blackett's dependence, the giromagnetic ratios of celestial bodies are approximately equal to $\sqrt{G}/c$. It has a simple explanation too. When there is the gravity induced electric polarization of a substance of a celestial body, its rotation must induce a magnetic field (Fig.(8)). It is important that all (composed by eN-plasma) celestial bodies - planets, stars, pulsars - obey the Blackett's dependence. It confirms a consideration that the gravity induced electric polarization must be characterizing for all kind of plasma. The calculation of magnetic fields of hot stars shows that they must be proportional to rotation velocity of stars (147). Magnetic fields of Ap-stars are measured, and they can be compared with periods of changing of luminosity of these stars. It is possible that this mechanism is characteristic for stars with rapid rotation (Fig.(9)), but obviously there are other unaccounted factors.

Taking into account the gravity induced electric polarization and coming from the Clairault's theory, we can describe the periastron rotation of binary stars as effect descended from non-spherical forms of star cores. It gives the quantitative explanation of this effect, which is in a good agreement with measuring data (Fig.(10)).

The solar oscillations can be considered as elastic vibrations of the solar core. It permits to obtain two basic frequencies of this oscillation: the basic frequency of sound radial oscillation of the core and the frequency of splitting depending on oscillations of substance density near its equilibrium value (Fig.(12)).

The plasma can exists in four possible states. The non-relativistic electron gas of plasma can be degenerate and non-degenerate. Plasma with relativistic electron gas can have a cold and a hot nuclear subsystem. Together with the atomic substance and neutron substance, it gives seven possible states. It suggests a way of a possible classification of celestial bodies. The advantage of this method of classification is in the possibility to estimate theoretically main parameters characterizing the celestial bodies of each class. And these predicted parameters are in agreement with astronomical observations. It can be supposed hypothetically that cosmologic transitions between these classes go in direction of their temperature being lowered. But these suppositions have no formal base

at all.

Discussing formulas obtained, which describe star properties, one can note an important moment: these considerations permit to look at this problem from different points of view. On the one hand, the series of conclusions follows from existence of spectrum of star mass (Fig.(1)) and from known chemical composition dependence. On the another hand, the calculation of natural frequencies of the solar core gives a different approach to a problem of chemical composition determination. It is important that for the Sun, both these approaches lead to the same conclusion independently and unambiguously. It gives a confidence in reliability of obtained results.

In fact, the calculation of magnetic fields of hot stars agrees with measuring data on order of the value only. But one must not expect the best agreement in this case because calculations were made for the case of a spherically symmetric model and measuring data are obtained for stars where this symmetry is obviously violated. But it is important, that the all remaining measuring data (all known of today) confirm both - the new postulate and formulas based on it. At that, the main stellar parameters - masses, radii and temperatures - are expressed through combinations of world constants and at that they show a good accordance with observation data. It is important that quite a satisfactory quantitative agreement of obtained results and measuring data can be achieved by simple and clear physical methods without use of any fitting parameter. It gives a special charm and attractiveness to star physics.

References

- [1] Allen C.W. - Astrophysical quantities, 1955, University of London, The Athlone Press.
- [2] Beskin V.S., Gurevich A.V., Istomin Ya.N.:
Physics of the Pulsar Magnetosphere (Cambridge University Press) (1993)
- [3] Blackett P.M.S.: *Nature*, **159**, 658, (1947)
- [4] Chandrasekhar S.: Monthly Notices of the RAS **93**, 449, (1933)
- [5] Christensen-Dalsgaard, J.: Stellar oscillation, Institut for Fysik og Astronomi, Aarhus Universitet, Denmark, (2003)
- [6] Elsworth, Y. et al. - In Proc. GONG'94 Helio- and Astero-seismology from Earth and Space, eds. Ulrich, R.K., Rhodes Jr, E.J. and Däppen, W., Astronomical Society of the Pacific Conference Series, vol. 76, San Francisco, **76**, 51-54.
- [7] Heintz W.D.: *Double stars* In Geoph. and Astroph. monographs, **15**, D.Reidel Publ. Corp., (1978)
- [8] Khaliullin K.F.: *Dissertation*, Sternberg Astronomical Institute, Moscow, (Russian) (2004) (see Table in Appendix on <http://astro07.narod.ru>)
- [9] Landau L.D. and Lifshits E.M.: Statistical Physics, **1**, 3rd edition, Oxford: Pergamon, (1980)
- [10] Landau L.D. and Lifshits E.M.: Electrodynamics of condensed matter, **1**, 3rd edition, Oxford: Pergamon, (1980)
- [11] Landau L.D. and Lifshits E.M.: The Classical Theory of Fields. **1**, Pergamon Press, N.Y. (1971)
- [12] Leung Y.C.: *Physics of Dense Matter* In Science Press/World Scientific, Beijing and Singapore, (1984)
- [13] I.I. Romanyuk et al. Magnetic Fields of Chemically Peculiar and Related Stars, Proceedings of the International Conference (Nizhnij Arkhyz, Special Astrophysical Observatory of Russian Academy of Sciences, September 24-27, 1999), eds: Yu. V. Glagolevskij and I.I. Romanyuk, Moscow, 2000, pp. 18-50.
- [14] Russel H.N.: Monthly Notices of the RAS **88**, 642, (1928)
- [15] Sirag S.-P.: *Nature*, **275**, 535, (1979)
- [16] Solar Physics, **175/2**, (<http://sohowww.nascom.nasa.gov/gallery/Helioseismology>)
- [17] Thorsett S.E. and Chakrabarty D.: *E-preprint: astro-ph/9803260*, (1998)

- [18] Vasiliev B.V. and Luboshits V.L.: *Physics-Uspekhi*, **37**, 345, (1994)
- [19] Vasiliev B.V.: *Nuovo Cimento B*, **116**, pp.617-634, (2001)

Appendix

The Table of main parameters of close binary stars
(cited on Kh.F.Khaliullin's dissertation, Sternberg Astronomical Institute. In Russian)

N	Name of star	U period of apsidal rotation, years	P period of ellipsoidal rotation, days	$M_1/M_\odot$ mass of component 1, in the Sun mass	$M_2/M_\odot$ mass of component 2, in the Sun mass	$R_1/R_\odot$ radius of 1 component in the Sun radius	$R_2/R_\odot$ radius of 2 component in the Sun radius	T_1 temperature of 1 component, K	T_2 temperature of 2 component, K	References
1	BW Aqr	5140	6.720	1.48	1.38	1.803	2.075	6100	6000	1,2
2	V 889 Aql	23200	11.121	2.40	2.20	2.028	1.826	9900	9400	3,4
3	V 539 Ara	150	3.169	6.24	5.31	4.512	3.425	17800	17000	5,12,24,67
4	AS Cam	2250	3.431	3.31	2.51	2.580	1.912	11500	10000	7,13
5	EM Car	42	3.414	22.80	21.40	9.350	8.348	33100	32400	8
6	GL Car	25	2.422	13.50	13.00	4.998	4.726	28800	28800	9
7	QX Car	361	4.478	9.27	8.48	4.292	4.054	23400	22400	10,11,12
8	AR Cas	922	6.066	6.70	1.90	4.591	1.808	18200	8700	14,15
9	IT Cas	404	3.897	1.40	1.40	1.616	1.644	6450	6400	84,85
10	OX Cas	40	2.489	7.20	6.30	4.690	4.543	23800	23000	16,17
11	PV Cas	91	1.750	2.79	2.79	2.264	2.264	11200	11200	18,19
12	KT Cen	260	4.130	5.30	5.00	4.028	3.745	16200	15800	20,21
13	V 346 Cen	321	6.322	11.80	8.40	8.263	4.190	23700	22400	20,22
14	CW Cep	45	2.729	11.60	11.10	5.392	4.954	26300	25700	23,24
15	EK Cep	4300	4.428	2.02	1.12	1.574	1.332	10000	6400	25,26,27,6
16	α Cr B	46000	17.360	2.58	0.92	3.314	0.955	9100	5400	28,29
17	Y Cyg	48	2.997	17.50	17.30	6.022	5.680	33100	32400	23,30
18	Y 380 Cyg	1550	12.426	14.30	8.00	17.080	4.300	20700	21600	31
19	V 453 Cyg	71	3.890	14.50	11.30	8.607	5.410	26600	26000	17,32,33
20	V 477 Cyg	351	2.347	1.79	1.35	1.567	1.269	8550	6500	34,35
21	V 478 Cyg	26	2.881	16.30	16.60	7.422	7.422	29800	29800	36,37
22	V 541 Cyg	40000	15.338	2.69	2.60	2.013	1.900	10900	10800	38,39
23	V 1143 Cyg	10300	7.641	1.39	1.35	1.440	1.226	6500	6400	40,41,42
24	V 1765 Cyg	1932	13.374	23.50	11.70	19.960	6.522	25700	25100	28
25	DI Her	29000	10.550	5.15	4.52	2.478	2.689	17000	15100	44,45,46,47
26	HS Her	92	1.637	4.25	1.49	2.709	1.485	15300	7700	48,49
27	CO Lac	44	1.542	3.13	2.75	2.533	2.128	11400	10900	50,51,52
28	GG Lup	101	1.850	4.12	2.51	2.644	1.917	14400	10500	17
29	RU Mon	348	3.585	3.60	3.33	2.554	2.291	12900	12600	54,55
30	GN Nor	500	5.703	2.50	2.50	4.591	4.591	7800	7800	56,57
31	U Oph	21	1.677	5.02	4.52	3.311	3.110	16400	15200	53,58,37
32	V 451 Oph	170	2.197	2.77	2.35	2.538	1.862	10900	9800	59,60
33	β Ori	228	5.732	19.80	7.50	14.160	8.072	26600	17800	61,62,63
34	FT Ori	481	3.150	2.50	2.30	1.890	1.799	10600	9500	64
35	AG Per	76	2.029	5.36	4.90	2.995	2.606	17000	17000	23,24
36	IQ Per	119	1.744	3.51	1.73	2.445	1.503	13300	8100	65,66
37	ζ Phe	44	1.670	3.93	2.55	2.851	1.852	14100	10500	11,67
38	KX Pup	170	2.147	2.50	1.80	2.333	1.593	10200	8100	21
39	NO Pup	37	1.257	2.88	1.50	2.028	1.419	11400	7000	11,69
40	VV Pyx	3200	4.596	2.10	2.10	2.167	2.167	8700	8700	70,71
41	YY Sgr	297	2.628	2.36	2.29	2.196	1.992	9300	9300	72
42	V 523 Sgr	203	2.324	2.10	1.90	2.682	1.839	8300	8300	73
43	V 526 Sgr	156	1.919	2.11	1.66	1.900	1.597	7600	7600	74
44	V 1647 Sgr	592	3.283	2.19	1.97	1.832	1.669	8900	8900	75
45	V 2283 Sgr	570	3.471	3.00	2.22	1.957	1.656	9800	9800	76,77
46	V 760 Sco	40	1.731	4.98	4.62	3.015	2.642	15800	15800	78
47	AO Vel	50	1.585	3.20	2.90	2.623	2.954	10700	10700	79
48	EO Vel	1600	5.330	3.21	2.77	3.145	3.284	10100	10100	21,63
49	α Vir	140	4.015	10.80	6.80	8.097	4.394	19000	19000	80,81,68
50	DR Vul	36	2.251	13.20	12.10	4.814	4.369	28000	28000	82,83

References

- [1] Khaliulilin Kh.F. and Kozyreva V.S.
Apsidal motion in the eclipsing binary system of BW Aqr
Astrophys. and Space Sci., **120** (1986) 9-16.
- [2] Imbert M.
Photoelectric radial velocities of eclipsing binaries. IV. Orbital elements of BW Aqr,
Astron.Astrophys.Suppl., **69** (1987) 397-401.
- [3] Khaliulilin Kh.F. and Khaliulilina A.I.
Fotometrisheskoe issledovanie zatmenno-dvoynoi sistemy s relativistskim vrasheniem orbity V889 Aql,
Astronom.zh.,**66**(1989)76-83 (in Russian).
- [4] Khaliulilin Kh.F. and Khaliulilina A.I.
K probleme vrashenia linii apsid v zatmennoi sisteme V889 Aql,
Astron.cirk., **N1486** (1987) 5-7 (in Russian).
- [5] Clausen J.V.
V 539 Arae: first accurate dimensions of eclipsing binaries,
Astron.Astrophys., **308** (1996) 151-169.
- [6] Lavrov M.I. and Lavrova N.V.
Revisia elementov fotometrisheskoi orbity EK Cep,
Astron.cirk. **971** (1977) 3-4 (in Russian).
- [7] Khaliulilin Kh.F. and Kozyreva V.S. *Apsidal motion in the eclipsing binary AS Cam*,
Astrophys. and Space Sci., **120** (1994) 115-122.
- [8] Andersen J. and Clausen J.V., *Absolute dimensions of eclipsing binaries.XV. EM Cainae*,
Astron.Astrophys. **213** (1989) 183-194.
- [9] Gemenez A. and Clausen J.V.,
Four-color photometry of eclipsing binaries. XXIIA. Photometric elements and apsidal motion of GL Cainae,
Astron.Astrophys. **161** (1986) 275-286.
- [10] Andersen J., Clausen J.V., Nordstrom B. and Reipurth B.,
Absolute dimensions of eclipsing binaries.I. The early-type detached system QX Cainae,
Astron.Astrophys. **121** (1983) 271-280.

- [11] Gemenez A., Clausen J.V. and Jensen K.S.
Four-color photometry of eclipsing binaries. XXIV. Aspidal motion of QX Cainae, ξ Phoenicis and NO Puppis,
Astron.Astrophys. **159** (1986) 157-165.
- [12] De Greve J.P.
Evolutionary models for detached close binaries: the system Arae and QX Cainae,
Astron.Astrophys. **213** (1989) 195-203.
- [13] Malony F.P., Guinan E.F. and Mukherjee J.
Eclipsing binary star as test of gravity theories
Astron.J. **102** (1991) 256-261.
- [14] Mossakovskaya L.V.
New photometric elements of AR Cas, an eclipsing binary system with apsidal motion
Astron. and Astroph. Trans. **2** (1992) 163-167.
- [15] Haffer C.M. and Collins G.M.
Computation of elements of eclipsing binary stars by high-speed computing machines
Astroph.J.Suppl., **7** (1962) 351-410.
- [16] Crinklaw G. and Etzel P.
A photometric analysis of the eclipsing binary OX Cassiopeiae
Astron.J. **98** (1989) 1418-1426.
- [17] Claret A. and Gimenez A.
The apsidal motion test of the internal stellar structure: comparison between theory and observations
Astron.Astroph. **277** (1993) 487-502.
- [18] Wolf M.
Aspidal motion in the eclipsing binary PV Cassiopeiae
Monthly Not.Roy.Soc. **286** (1995) 875-878.
- [19] Popper D.M.
Rediscussion of eclipsing binaries.XVII.The detached early A type binaries PV Cassiopeae and WX Cephei
Astron.J. **93** (1987) 672-677.

- [20] Lavrov M.I. and Lavrova N.V.
Revisia fotometricheskih elementov u zatmennykh dvoynykh sistem s ekscentricheskimi orbitami.2.KT Cen
Trudy Kaz.Gor.AO **49** (1985) 18-24 (in Russian).
- [21] Soderhjelm S.
Observations of six southern eclipsing binaries for apsidal motion
Astron.Astroph.Suppl.Ser **22** (1975) 263-283.
- [22] Gemenez A., Clausen J.V. and Anderson J.
Four-color photometry of eclipsing binaries. XXIA. Photometric analysis and apsidal motion study of V346 Centauri,
Astron.Astrophys. **160** (1986) 310-320.
- [23] Gemenez A., Chun-Hwey Kim and Il-Seong Nha
Apsidal motion in the early-type eclipsing binaries CW Cephei, Y Cyg and AG Per
Montly .Not.Roy.Astron.Soc. **224** (1987) 543-555.
- [24] Bocula R.A.
Peresmotr elementov fotometricheskoi orbity zatmennykh sistem CW Cep, V 539 Ara, AG Per, AR Aur, RS Cha, ZZ Boo.
Peremennye zvezdy**21** (1983) 851-859 (in Russian).
- [25] Khaliulilin Kh.F.
Relativistskoe vrashenie orbity zatmennoi dvoynoi sistemy EK Cep
Astron.zh.**60** (1983) 72-82 (in Russian).
- [26] Tomkin J.
Secondaries of eclipsing binary. V. EK Cephei
Astroph.J. **271** (1983) 717-724.
- [27] Claret A., Gemenez A. and Martin E.L.
A test case of stellar evolution the eclipsing binary EK Cephei
Astron.Astroph. **302** (1995) 741-744.
- [28] Volkov I.M.
The discovery of apsidal motion in the eclipsing binary system α Cr B
Inf.Bull.Var.Stars N3876,(1993) 1-2.
- [29] Quiroga R.J., van L.P.R.
Angular momenta in binary systems
Astroph.Space Sci. **146** (1988) 99-137.

- [30] Hill G. and Holmgren D.E.
Studies of early-type variable stars
Astron.Astroph.**297** (1995) 127-134.
- [31] Hill G. and Batten A.H.
Studies of early-type variable stars.III. The orbit and physical dimensions for V 380 Cygni
Astron.Astroph.**141** (1984) 39-48.
- [32] Zakirov M.M.
Ob apsidalnom dvizhenii v dvoinoi sisteme V 453 Cyg
Astron.cirk.N1537,21 (in Russian).
- [33] Karetnikov V.G.
Spectral investigation of eclipsing binary stars at the stage of mass exchange
Publ.Astron.Inst.Czech.**70** (1987) 273-276.
- [34] Mossakovskaya L.V. and Khaliulilin Kh.F.
Prichina anomal'nogo apsidalnogo dvizhenia v sisteme V 477 Cyg
Astron.cirk.N1536, 23-24 (in Russian).
- [35] Gemenez A. and Quintana J.M.
Apsidal motion and revised photometry elements of the eccentric eclipsing binary V 477 Cyg
Astron.Astrophys. **260** (1992) 227-236.
- [36] Mossakovskaya L.V. and Khaliulilin Kh.F.
Vrashenie linii apsid v sisteme V 478 Cyg
Pisma v Astron.zh.**22** (1996) 149-152.
- [37] Popper D.M. and Hill G.
Rediscussion of eclipsing binaries.XVII.Spectroscopic orbit of OB system with a cross-correlation procedure
Astron.J. **101** (1991) 600-615.
- [38] Khaliulilin Kh.F.
The unique eclipsing binary system V 541 Cygni with relativistic apsidal motion
Astrophys.J. **229** (1985) 668-673.
- [39] Lines R.D., Lines H., Guinan E.F. and Carroll
Time of minimum determination of eclipsing binary V 541 Cygni
Inf.Bull.Var.Stars N3286,1-3.

- [40] Khaliulilin Kh.F.
Vrashenie linii apsid v zatmennoi sisteme V 1143 Cyg
Asrton. cirk.N1262,1-3 (in Russian).
- [41] Andersen J., Garcia J.M., Gimenes A. and Nordstrom B.
Absolute dimension of eclipsing binaries.X. V1143 Cyg
Astron.Astrophys. **174** (1987) 107-115.
- [42] Burns J.F., Guinan E.F. and Marshall J.J.
New apsidal motion determination of eccentric eclipsing binary V 1143 Cyg
Inf.Bull.Var.Stars N4363,1-4.
- [43] Hill G. and Fisher W.A.
Studies of early-type variable stars.II. The orbit and masses of HR 7551
Astron.Astrophys. **139** (1985) 123-131.
- [44] Martynov D.Ya. and Khaliulilin Kh.F.
On the relativistic motion of periastron in the eclipsing binary system DI Her
Astrophys.and Space Sci. **71** (1980) 147-170.
- [45] Popper D.M.
Rediscussion of eclipsing binaries.XVII. DI Herculis, a B-type system with an eccentric orbit
Astron.J. **254** (1982) 203-213.
- [46] Martynov D.Ia. Lavrov M.I.
Revizia elementov fotometricheskoi orbity i skorosti vrashenia linii apsid u zatmennoi dvoimoi sistemy DI Her
Pisma v Astron.Zh. **13** (1987) 218-222 (in Russian).
- [47] Khaliulilin Kh.F., Khodykin S.A. and Zakharov A.I.
On the nature of the anomalously slow apsidal motion of DI Herculis
Astrophys.J. **375** (1991) 314-320.
- [48] Khaliulilina A.I. and Khaliulilin Kh.F.
Vrashenie linii apsid v zatmennoi dvoimoi sisteme HS Her
Astron.cirk.N 1552 (1992) 15-16(in Russian).
- [49] Martynov D.Ia., Voloshina I.E. and Khaliulilina A.I.
Fotometricheskie elementy zatmennoi sistemy HS Her
Asrton. zh. **65** (1988) 1225-1229 (in Russian).

- [50] Mezzetti M., Predolin F., Giuricin G. and Mardirossian F.
Revised photometric elements of eight eclipsing binaries
Astron.Astroph.Suppl. **42** (1980) 15-22.
- [51] Mossakovskaya L.V. and Khaliulilin Kh.F.
Tret'e telo v zatmennoi sisteme s apsidalnym dvizheniem CO Lac?
Astron. cirk.N1495, 5-6 (in Russian).
- [52] Semeniuk I.
Apsidal motion in binary systems. I. CO Lacertae, an eclipsing variable with apsidal motion
Acta Astron. **17** (1967) 223-224.
- [53] Andersen J.
Accurate masses and radii of normal stars
Astron.Astroph.Rev. **3** (1991) 91-126.
- [54] Khaliulilina A.I., Khaliulilin Kh.F. and Martynov D.Ya.
Apsidal motion and the third body in the system RU Monocerotis
Montly .Not.Roy.Astron.Soc. **216** (1985) 909-922.
- [55] Martynov D.Ya. and Khaliulilina A.I.
RU Monocerotis: poslednie resultaty
Astron.zh.**63** (1986) 288-297 (in Russian).
- [56] Shneller H.
Uber die periodenanderrungen bei bedeckungsveranderlichen
Budd.Mitt. N53 (1962) 3-41.
- [57] Kort S.J., J. de, *The orbit and motion of priastron of GN Normae*
Ricerche Astron. **3** (1954) 119-128.
- [58] Kamper B.C.
Light-time orbit and apsidal motion of eclipsing binary U Ophiuchi
Astrophys. Space Sci. **120** (1986) 167-189.
- [59] Clausen J.V., Gemenez A. and Scarfe C.
Absolute dimentions of eclipsing binaries.XI. V 451 Ophiuchi
Astron.Astroph. **167** (1986) 287-296.
- [60] Khaliulilin Kh.F. and Kozyreva V.S.
Photometric light curves and physical parameters of eclipsing binary systems IT Cas, CO Cep, AI Hya with possible apsidal motions
Astrophys. and Space Sci., **155** (1989) 53-69.

- [61] Monet D.G.
A discussion of apsidal motion detected in selected spectroscopic binary systems
 Astrophys. J., **237** (1980) 513-528.
- [62] Svechnikov M.A.
Katalog orbitalnyh elementov, mass i svetimostei tesnyh dvoinyh zvezd
 Irkutsk, Izd-vo Irkutsk. Univer.(In Russian).
- [63] Brancewicz H.K. and Dworak T.Z.
A Catalogue of parameters for eclipsing binaries
 Acta Astron., **30** (1980) 501-524.
- [64] Wolf M. and Saronova L.,
Apsidal motion in the eclipsing binary FT Ori
 Astron.Astroph. Suppl.**114** (1995) 143-146.
- [65] Drozd M., Krzesinski J. and Paydosz G.,
Apsidal motion of IQ Persei
 Inf. Bull.Var.Stars, N3494, 1-4.
- [66] Lacy C.H.S. and Fruch M.L.
Absolute dimentions and masses of eclipsing binaries. V. IQ Persei
 Astroph.J.**295** (1985) 569-579.
- [67] Andersen J.
Spectroscopic observations of eclipsing binaries.V. Accurate mass determination for the B-type systems V 539 Arae and ξ Phaeniceis
 Astron.Astroph.**118** (1983) 255-261.
- [68] Odell A.P.
The structure of Alpha Virginis.II. The apsidal constant
 Astroph.J.**192** (1974) 417-424.
- [69] Gronbech B.
Four-color photometry of eclipsing binaries.V. photometric elements of NO Puppis
 Astron.Astroph.**50** (1980) 79-84.
- [70] Harmanec P.
Stellar masses and radii based on motion binary data
 Bull.Astron.Inst.Czech.**39** (1988) 329-345.

- [71] Andersen J., Clausen L.V. and Nordstrom B.
Absolute dinemtions of eclipsing binaries.V. VV Pyxidis a detached early A-tipe system with equal components
Astron.Astroph.**134** (1984) 147-157.
- [72] Lacy C.H.S.
The photometric orbit and apsidal motion of YY Sagittarii
Astroph.J.**105** (1993) 637-645.
- [73] Lacy C.H.S.
The photometric orbit and apsidal motion of V 523 Sagittarii
Astroph.J.**105** (1993) 630-636.
- [74] Lacy C.H.S.
The photometric orbit and apsidal motion of V 526 Sagittarii
Astroph.J.**105** (1993) 1096-1102.
- [75] Andersen J. and Gimenes A.
Absolute dinemtions of eclipsing binaries.VII. V 1647 Sagittarii
Astron.Astroph.**145** (1985) 206-214.
- [76] Swope H.H.
V 2283 Sgr, an eclipsing star with rotating apse
Ric.Astron.**8** (1974) 481-490.
- [77] O’Konnell D.J.K.
The photometric orbit and apsidal motion of V2283 Sagittarii
Ric.Astron.**8** (1974) 491-497.
- [78] Andersen J., Clausen L.V., Nordstrom B. and Popper D.M.
Absolute dinemtions of eclipsing binaries.VIII. V 760 Scorpii
Astron.Astroph.**151** (1985) 329-339.
- [79] Clausen L.V., Gimenez A. and Houten C.J.
Four-color photometry of eclipsing binaries.XXVII. A photometric anallysis of the (possible) Ap system AO Velorum
Astron.Astroph.**302** (1995) 79-84.
- [80] Popper D.M.
Stellar masses
Ann. Rev. Astron. and Astroph.**18** (1980) 115-164.

- [81] Dukes R.J.
The beta Cephei nature of Spica
 Astroph.J. **192** (1974) 81-91.
- [82] Khaliulilina A.I.
DR Vulpeculae: the quadruple system
 Montly .Not.Roy.Astron.Soc. **225** (1987) 425-436.
- [83] Khaliulilin Kh.F. and Khaliulilina A.I.
Fotometrisheskoe issledovanie zatmennoi zvezdy DR Vul. Parametry sistemy i vrashenie linii apsid,
 Astron.zh., **N65** (1988) 108-116 (in Russian).
- [84] Khaliulilin Kh.F. and Kozyreva V.S.
Photometric light curves and physical parameters of eclipsing binary systems IT Cas, CO Cep, AI Hya with possible apsidal motions
 Astrophys. and Space Sci., **155** (1989) 53-69.
- [85] Holmgren D. and Wolf M. *Apsidal motion of the eclipsing binary IT Cassiopeiae*
 Observatory **116** (1996) 307-312.